

Burnout among chronic pain anesthesiologists

Alaa Abd-Elseyed

ABSTRACT

Burnout is a concern to individuals in different professions and it continues to be a major concern among physicians in different specialties. Hyman *et al*¹ conducted a study to compare the severity of burnout among different anesthesiology subspecialties. Chronic pain anesthesiologists had higher burnout when compared with other surveyed subspecialties.

Anesthesiology, University of Wisconsin Madison School of Medicine and Public Health, Madison, Wisconsin, USA

Correspondence to Dr Alaa Abd-Elseyed, University of Wisconsin Madison School of Medicine and Public Health, Madison, WI 53726, USA; alaaawny@hotmail.com

Correction notice This article has been corrected since it published Online First. The infographic has been updated.

Contributors AA-E designed and prepared the infographic.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© American Society of Regional Anesthesia & Pain Medicine 2021. No commercial re-use. See rights and permissions. Published by BMJ.




To cite Abd-Elseyed A. *Reg Anesth Pain Med* 2021;**46**:388.

Received 25 January 2021
Accepted 26 January 2021
Published Online First 11 February 2021

Regional Anesthesia & Pain Medicine

Burnout Among Chronic Pain Anesthesiologists



doi:10.1136/rapm-2021-102530



► <http://dx.doi.org/10.1136/rapm-2020-101520>
Reg Anesth Pain Med 2021;**46**:388.

REFERENCE

- Hyman S, Card E, de Leon-Casasola O, *et al*. Prevalence of burnout and its relationship to health status and social support in more than 1,000 subspecialty Anesthesiologists 2021;**46**:381–7.