# Optimal location for continuous catheter analgesia among the femoral triangle, proximal, or distal adductor canal after total knee arthroplasty: a randomized double-blind controlled trial

Bora Lee,<sup>1</sup> Sang Jun Park,<sup>1</sup> Kwan Kyu Park,<sup>2</sup> Hee Jung Kim,<sup>1</sup> Yong Suk Lee,<sup>1</sup> Yong Seon Choi<sup>1</sup>

# ABSTRACT

► Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi.org/ 10.1136/rapm-2021-103284).

<sup>1</sup>Department of Anesthesiology and Pain Medicine, Anesthesia and Pain Research Institute, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea <sup>2</sup>Department of Orthopedic Surgery, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea

#### Correspondence to

Dr Yong Seon Choi, Department of Anesthesiology and Pain Medicine, Yonsei University College of Medicine, Seoul, Republic of Korea; YSCHOI@yuhs.ac

Received 27 October 2021 Accepted 7 February 2022 Published Online First 16 February 2022 Background Pain management after total knee arthroplasty is essential to improve early mobilization, rehabilitation, and recovery. Continuous adductor canal (AC) block provides postoperative analgesia while preserving guadriceps strength. However, there have been inconsistencies regarding the optimal location for continuous catheter block. We compared continuous femoral triangle, proximal AC, and distal AC blocks for postoperative analgesia after total knee arthroplasty. Methods Patients undergoing unilateral total knee arthroplasty were randomly assigned to three groups: femoral triangle, proximal AC, or distal AC. The surgeon performed periarticular local anesthetic infiltration. After surgery, an ultrasound-quided perineural catheter insertion procedure was performed. The primary endpoint was pain scores at rest in the morning on the first postoperative day. Secondary endpoints included pain scores at rest and during activity at other time points, guadriceps strength, and opioid consumption.

**Results** Ninety-five patients, 32 in the femoral triangle group, 31 in the proximal AC group, and 32 in the distal AC group, completed the study. Analysis of the primary outcome showed no significant difference in pain scores among groups. Secondary outcomes showed significantly lower pain scores at rest and during activity in the distal AC group than in the femoral triangle and proximal AC groups in the morning of the second postoperative day. Quadriceps strength and opioid consumption did not differ among groups.

**Conclusions** Continuous femoral triangle, proximal AC, and distal AC blocks in the setting of periarticular local anesthetic infiltration provide comparable postoperative analgesia after total knee arthroplasty.

Trial registration number NCT04206150.

**INTRODUCTION** 

#### Check for updates

© American Society of Regional Anesthesia & Pain Medicine 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Lee B, Park SJ, Park KK, et al. Reg Anesth Pain Med 2022;47:353–358.

(ACB) provides postoperative analgesia while

preserving quadriceps strength.<sup>3</sup>

Total knee arthroplasty (TKA) is associated with

moderate to severe pain in 50% of patients in

the first three postoperative days.<sup>1</sup> Optimal pain

management after TKA is therefore essential to

The adductor canal (AC) is a triangular musculoaponeurotic conduit between the apex of the femoral triangle and adductor hiatus. Previous clinical studies on continuous ACBs used surface anatomical landmarks to identify the proximal end of the AC as the midpoint between the anterior superior iliac spine (ASIS) and the superior border of the patella.<sup>4 5</sup> Recent cadaveric studies have revealed that the proximal end of the AC is located distal from the midpoint of the thigh.<sup>6-8</sup> In recent clinical studies regarding ACB catheter placement, the proximal end of the AC was precisely identified using ultrasound landmarks as the intersection of the medial borders of the sartorius muscle and the adductor longus muscle.<sup>9-11</sup> However, despite previous studies, the ideal location of continuous ACB catheter analgesia after TKA has been difficult to determine due to inconsistent AC identification and various catheter insertion positions.<sup>4 5 9-11</sup> In addition, most studies have compared femoral triangle block with ACB for continuous catheter analgesia, showing inconsistent results.<sup>5 9 10</sup> Therefore, a comparison of analgesic efficacy provided by catheters inserted into the femoral triangle, proximal AC, and distal AC may provide an objective basis for a clear recommendation on this approach.

We designed this randomized, controlled, doubleblind clinical trial to compare the effects of continuous femoral triangle, proximal AC, and distal AC blocks on postoperative analgesia after TKA. Our primary endpoint was to compare pain scores in the morning on postoperative day 1 (POD1). The secondary endpoints included pain scores at other time points, quadriceps muscle strength, and opioid consumption.

# **METHODS**

This randomized, controlled, double-blind study was registered at ClinicalTrials.gov on December 20, 2019. A total of 96 adult patients with an American Society of Anesthesiologists physical status of class I–III scheduled for elective, unilateral, primary TKA were enrolled between April 2020 and July 2021. Exclusion criteria were chronic opioid use (daily use for >1 month of 20 mg oral morphine equivalents), revision surgery, contraindications to spinal anesthesia or peripheral nerve blocks, and allergy to lidocaine or ropivacaine. Written informed consent was obtained from all patients.

# **Original research**

Patients were randomly assigned to either the femoral triangle, proximal AC, or distal AC groups according to a computergenerated randomization sequence on the day of surgery. Patient allocation was performed by an investigator who was not involved in postoperative assessments. Randomization assignments were concealed until catheter insertion. Aside from the investigator performing the catheter insertion procedure, other investigators, surgeons, patients, attending anesthesiologists, and nurses were blinded to group assignments during the study period.

#### Anesthesia and surgery protocol

On the patients' arrival at the operating room, routine monitoring was performed. The patients received spinal anesthesia with 10-12 mg of 0.5% hyperbaric bupivacaine. All TKAs were performed via the medial parapatellar approach by a single surgical team. The surgeon performed periarticular local anesthetic infiltration with 150 mg of ropivacaine and 30 mg of ketorolac diluted with normal saline to a total volume of 50 mL. The first 20 mL of the mixture were injected into the posterior capsule, the medial and lateral collateral ligaments, and medial and lateral meniscus remnants, prior to implant placement. The remaining 30 mL were used to infiltrate the soft tissue around the knee, including the surrounding muscles, synovium, fat pad, and subcutaneous tissue, before wound closure. All patients received preoperative celecoxib (200 mg) orally and intraoperative doses of acetaminophen (1 g), tranexamic acid (1 g), and dexamethasone (5 mg) intravenously. In the ward, all patients received celecoxib 200 mg orally and acetaminophen 1 g intravenously every 12 hours thereafter. Intravenous tramadol (50 mg) was administered as rescue analgesia. Patients received standard of care physical therapy sessions (two times on POD1 and POD2) and were mobilized once on the day of surgery with caregiver assistance and ambulated as soon as possible on POD1.

#### **Catheter insertion procedure**

After surgery, the patient was transferred to a block room and routine monitoring was performed. All catheters were inserted under a full aseptic technique using a real-time ultrasoundguided in-plane approach by an experienced anesthesiologist. A multi-hole perineural catheter through a catheter-over-needle system (E-cath PLUS, PAJUNK GmbH, Geisingen, Germany) and a linear 6–13-MHz ultrasound probe (HFL38xp, SonoSite Inc., Bothell, Washington, USA) were used for all patients.

On ultrasound, the proximal end of the AC was defined as the intersection of the medial borders of the sartorius muscle and the adductor longus muscle (online supplemental figure 1). In the femoral triangle group, the insertion site was determined at 2/15th the femur length above the proximal end of the AC along the long axis of the femur (figure 1A). In the proximal AC group, the insertion site was the proximal end of the AC (figure 1B). In the distal AC group, the insertion site was determined at 1/15th the femur length below the proximal end of the AC along the long axis of the femur (figure 1C). In our hospital's TKA protocol, a full-length standing anteroposterior radiograph is obtained before admission. Based on this, femur length was measured from the top of the femoral head to the most distal end of the medial femoral condyle (online supplemental figure 2).<sup>12</sup>

In preliminary data from 10 patients, the mid-thigh was approximated at 2/15th of the femur length above the proximal end of the AC (online supplemental table 1). In addition, an area 2 cm above the adductor hiatus was approximated at 1/15th of the femur length below the proximal end of the AC.



**Figure 1** Location of catheter insertion and corresponding ultrasound images. (A) The probe moves cranially along the long axis of the thigh as 2/15th of the femur length from the proximal end of the AC. (B) The probe is positioned to identify the proximal end of the AC, where the medial borders of ALM and the sartorius muscle intersect. (C) The probe moves caudally along the long axis of the thigh as 1/15th of the femur length from the proximal end of the AC. AC, adductor canal; ALM, adductor longus muscle; AMM, adductor magnus muscle; FA, femoral artery; FV, femoral vein; SN, saphenous nerve; VMM, vastus medialis muscle.

The proximal end of the AC was identified by ultrasound and the length from the ASIS to the superior border of the patella was measured using a tape measure.

Following identification of the designated site, an 18-gage cannula with an indwelling 21-gage needle was advanced through the sartorius muscle with the needle tip positioned lateral to the saphenous nerve. To open a space for catheter insertion, 10 mL of 0.2% ropivacaine was injected via the needle for hydrodissection. A 21-gage multi-hole E-catheter was inserted through an indwelling 18-gage cannula. An additional 5 mL of 0.2% ropivacaine was injected during ultrasound imaging to ensure correct placement of the catheter tip between the artery and the deep fascia of the sartorius in the femoral triangle group or between the artery and the vastoadductor membrane (VAM) in the AC groups (online supplemental figure 3). The catheter was secured with a sterile occlusive dressing and an anchoring device. During the 48-hour postoperative period, 0.2% ropivacaine was infused via the perineural catheter at a basal rate of 6 mL/hour, a 4 mL bolus, and a lockout time of 30 min using a portable, patientcontrolled infusion pump. After the procedure, all patients were dressed with hospital uniform tops and bottoms to blind the group assignment.

## **Outcome measurement**

The primary endpoint was pain scores at rest in the morning on POD1. The pain intensity at rest and during activity was evaluated using an 11-point numeric rating scale (NRS: 0=no pain, 10=worst imaginable pain). The secondary endpoints included pain scores at other time points, quadriceps muscle strength, opioid consumption, and local anesthetic consumption. Pain scores and quadriceps strength were assessed at four-time points: preoperative baseline, the morning (08:00-09:00) of POD1, the afternoon (16:00-17:00) of POD1, and the morning (08:00-09:00) of POD2. The quadriceps strength of both legs was tested using a hand-held dynamometer (Lafayette Manual Muscle Test System, Lafavette Instrument Company, Lafavette, Indiana, USA).<sup>13</sup> Patients were instructed to extend their knees two times, with a 30 s pause between each attempt. The maximum force achieved was used in the analysis. The consumption of tramadol was converted to oral morphine equivalents.<sup>14</sup> Preoperative data collection included the Hospital Anxiety and Depression Scale (HADS) ranging from 0 to 21. The probable presence of anxiety or depression was defined as a HADS anxiety or depression subscale score  $\geq 11.^{15}$ 

# **Statistical analysis**

We calculated the sample size required to detect a difference in NRS>3 among the three groups using pairwise comparison.<sup>5</sup> Bonferroni correction was performed to adjust for the increase in type I error in multiple comparisons. Accordingly, 29 subjects were required in each group to achieve a statistical power of 90% at p value of <0.0167. Therefore, 32 patients were enrolled in each group to account for a 10% dropout rate. The Shapiro-Wilk and Kolmogorov-Smirnov tests were used to confirm normality of the data distribution. One-way analysis of variance and the Kruskal-Wallis test were used to analyze parametric and nonparametric continuous variables, respectively. Post-hoc analysis was performed using the t-test or Mann-Whitney test as appropriate and adjusted using the Bonferroni correction. Intergroup comparisons of categorical variables were conducted using Fisher's exact test or the  $\chi^2$  test as appropriate. Continuous variables are presented as mean±SD or median (IQR); categorical variables are presented as numbers (percentages). Statistical analyses were performed using R V.3.5.1 (R Foundation for Statistical Computing, Vienna, Austria), SPSS V.23.0, (IBM Corp., Armonk, New York, USA), or MedCalc Statistical Software V.18.11.3 (MedCalc Software Ltd., Ostend, Belgium). Statistical significance (p) was set at < 0.05.

# RESULTS

Of the 102 patients screened for eligibility, 96 were enrolled in the study and randomly allocated to three groups: femoral triangle, proximal AC, or distal AC. One patient randomized to the proximal AC group experienced postoperative delirium (a disturbance in attention and awareness) and was withdrawn from the study during the postoperative follow-up period. Data from 95 patients were analyzed and patient flow through the study is shown in figure 2.

Patient characteristics, operative data, and baseline assessments were similar among groups (table 1), as was the probable presence of depression or anxiety (p=0.157 and p=0.189, respectively). Leg length measurement based on radiograph showed no differences among groups. In the femoral triangle group, catheters were inserted  $5.7 \pm 0.4$  cm proximally from the proximal end of the AC. In the distal AC group, catheters were inserted  $2.8 \pm 0.2$  cm distally from the proximal end of the AC.

Postoperative pain scores are shown in figure 3. There was no significant difference in NRS pain scores at rest and during activity among the three groups in the morning and afternoon



**Figure 2** Consolidated Standards of Reporting Trials study flow diagram. AC, adductor canal.

of POD1. NRS pain scores at rest (p=0.007) and during activity (p=0.005) were lower in the distal AC group in the morning of POD2 than those in the femoral triangle and proximal AC groups.

Dynamometer readings and analgesia are shown in table 2. Quadriceps strength measurements in the operative and nonoperative leg were not significantly different among groups in the morning and afternoon of POD1, and morning of POD2, respectively. Quadriceps strength measurements in the operated leg decreased from preoperative values in all groups at each time point. The cumulative amounts of 0.2% ropivacaine infused until 24, 36, and 48 hours postoperatively were comparable among groups. The number of patients who received rescue analgesics during the first 24 hours postoperatively was higher in the femoral triangle group than in the proximal and distal AC groups (p=0.027). However, cumulative opioid consumption was not significantly different among groups. There were no continuous block-related complications, such as falls, local anesthetic toxicity, or infection.

## DISCUSSION

In this study, we found no difference in postoperative pain scores at rest and during activity in the morning and afternoon of POD1 after TKA among continuous femoral triangle, proximal AC, and distal AC blocks in the setting of periarticular local anesthetic infiltration. Among secondary outcomes, postoperative pain scores at rest and during activity in the morning of POD2 were significantly reduced in continuous distal ACB compared with continuous femoral triangle block and continuous proximal ACB.

The ideal location of continuous ACB after TKA may provide prolonged pain relief and improved functional recovery while preserving muscle strength. However, there are no clear recommendations for this approach due to inconsistent AC identification and different catheter insertion positions.<sup>4 5 9-11</sup> Three studies showed no differences in analgesia between continuous femoral triangle block and ACB,<sup>4 9 10</sup> whereas two others showed better analgesia or opioid-sparing effect in a proximal location than a distal location.<sup>5</sup><sup>11</sup> In one study, pain score was lower on POD1 in the proximal ACB (midpoint of the thigh) than in the distal ACB (2-3 cm proximal to the adductor hiatus).<sup>5</sup> In another, cumulative sufentanil consumption within 24 hours after TKA was lower in the proximal ACB (proximal end of AC) than in the middle ACB (3-5 cm distal to the proximal end of AC).<sup>11</sup> Considering the most studied catheter positions,<sup>4 5 9–11</sup> we selected three catheter insertion locations for continuous catheter analgesia: femoral triangle, proximal AC, and distal AC. In

Table 1 Demographic data and baseline assessments						
	Femoral triangle group (N=32)	Proximal AC group (N=31)	Distal AC group (N=32)	P value		
Demographic data						
Age (years)	72.3±4.7	71.5±5.0	71.2±5.6	0.793		
Female/male	22 (69)	23 (74)	24 (75)	0.831		
Height (cm)	156.5±7.8	157.1±8.1	156.4±6.3	0.723		
Weight (kg)	65.5±8.4	65.5±9.8	63.9±7.5	0.474		
Body mass index (kg/m <sup>2</sup> )	26.8±3.2	26.4±2.4	26.1±2.7	0.667		
ASA class (I/II/III)	5/22/5	2/23/6	2/26/4	0.603		
Operation side						
Left/right	15/17	15/16	16/16	0.969		
Length of surgery (min)	47.5 (42.0–61.5)	47.0 (44.0–58.0)	49.5 (43.0–55.5)	0.998		
Baseline assessments						
HADS (0–21)						
Depression	8.7±3.9	8.5±4.4	7.8±3.2	0.452		
Anxiety	6.0 (4.0–10.0)	8.0 (3.0–9.0)	6.0 (4.0-8.0)	0.794		
NRS resting pain scores	2.0 (0–3.5)	2.0 (0–3.0)	2.0 (0–2.5)	0.529		
Baseline quadriceps strength (kgf)						
Operative leg	10.7±3.7	11.2±3.6	10.2±2.5	0.254		
Non-operative leg	12.6±3.6	13±3	12±2.7	0.251		
Femur length (cm)	42.8±2.8	42.6±3.2	42.2±2.6	0.623		

Values are presented as median (IQR), mean±SD, or number of patients (%).

AC, adductor canal; ASA, American Society of Anesthesiologists; HADS, Hospital Anxiety and Depression Scale; kgf, kilogram-force unit; NRS, numeric rating scale.





contrast to previous studies on ACBs,<sup>5</sup><sup>11</sup> catheter insertion was performed postoperatively to minimize catheter tip displacement due to knee movement during surgery without tunneling technique. Our methodology is unique in the determination of each catheter insertion location in relation to individual femur length based on both ultrasound and radiographic images. This has the advantage of being an objective measurement compared with only ultrasound images or surface anatomy.

Our primary outcome showed comparable postoperative analgesia after TKA with continuous femoral triangle, proximal AC, and distal AC blocks. This finding is consistent with most studies on continuous catheter analgesia after TKA.4 9 10 Our results could be explained by the musculoaponeurotic tunnel feature of the AC and the target nerves of local anesthetic administration. The spread of local anesthetic is determined by the fascial limits and the muscles surrounding the space.<sup>16</sup><sup>17</sup> The roof of the AC is a continuous fascia with thin proximal and thick distal portions.<sup>16</sup><sup>18</sup> The thin superficial layer is the VAM, which connects the medial edge of the vastus medialis to the lateral edge of the adductor magnus, and the thick deep layer is the aponeurosis of the vastus medialis oblique.<sup>16</sup> Because the femoral triangle is a space without VAM, longitudinal spread of local anesthetics may be less extensive than in AC.<sup>19</sup> Moreover, the two thick layers of the roof in the narrow distal AC space may enhance local anesthetic spread as they are histologically aponeurotic and stiff.<sup>16</sup> In a cadaveric study, 20 mL of dye injected into the distal femoral triangle stained the saphenous nerve and the nerve to vastus medialis in all specimens<sup>20</sup>; 10 mL of dye injected into the proximal end of the AC had spread to the saphenous nerve, posteromedial branch of the nerve to vastus medialis, superior medial genicular nerve, and genicular branch of the obturator nerve in all specimens.<sup>16</sup> When 10 mL of dye was injected into an area 1-2 cm proximal to the adductor hiatus had spread into the popliteal fossa and stained the popliteal plexus, genicular branch of the posterior obturator nerve, as well as the saphenous nerve.<sup>21</sup> However, 10–20 mL dye injected

Table 2 Quadriceps strength, perine	eural infusion pump data, and opio	id consumption		
	Femoral triangle group (N=32)	Proximal AC group (N=31)	Distal AC group (N=32)	P value
Quadriceps strength of operative leg (kgf)				
POD1 morning	5.0±2.8	5.0±2.7	4.9±2.9	0.885
POD1 afternoon	4.5±2.3	4.1±2.5	4.8±2.7	0.275
POD2 morning	4.2±3.0	4.8±2.4	5.9±2.3	0.274
Quadriceps strength of non-operative leg (kg	f)			
POD1 morning	11.7±4.0	12.1±3.1	10.6±3.6	0.125
POD1 afternoon	11.4±3.8	11.4±3.5	10.3±3.0	0.253
POD2 morning	11.5±3.5	12.2±3.3	11.1±3.3	0.346
Amount of 0.2% ropivacaine given as boluses	s plus background infusion (mL)			
0–24 hours	173.4 (159.7–187.1)	175.4 (157.7–193.1)	163.6 (155.7–185.2)	0.583
24–36 hours	83.8 (77.8–95.5)	87.7 (75.9–105.3)	81.9 (76.0–93.5)	0.556
36–48 hours	79.8 (73.8–83.8)	79.9 (77.9–85.8)	79.8 (72.0–88.0)	0.635
Amount of 0.2% ropivacaine given as boluses	s (mL)			
0–24 hours	26.0 (12.0–40.0)	28.0 (10.0–44.0)	16.0 (8.0–40.0)	0.573
24–36 hours	12.0 (8.0–24.0)	16.0 (4.0–34.0)	10.0 (4.0–22.0)	0.485
36–48 hours	8.0 (2.0–12.0)	8.0 (6.0–14.0)	8.0 (0.0–16.0)	0.764
Patients receiving rescue analgesics (n)				
0–24 hours	22 (68.8%)	12 (38.7%)	13 (40.6%)	0.027
24–36 hours	13 (40.6%)	12 (38.7%)	7 (21.9%)	0.219
36–48 hours	12 (37.5%)	8 (25.8%)	5 (15.6%)	0.138
Cumulative opioid consumption (morphine ed	quivalents)			
~24 hours	0.4 (0.0–0.4)	0.0 (0.0–0.3)	0.0 (0.0–0.4)	0.055
~36 hours	0.4 (0.0–0.8)	0.2 (0.0–0.4)	0.1 (0.0–0.4)	0.169
~48 hours	0.6 (0.1–1.0)	0.4 (0.0–0.5)	0.1 (0.0–0.5)	0.062

Values are presented as median (IQR), mean±SD, or number of patients (%).

\_AC, adductor canal; kgf, kilogram-force unit; POD, postoperative day.

near the adductor hiatus led to varying staining results of the nerve to vastus medialis (0%–30%) because the nerve to vastus medialis is separated from the structures within the proximal AC by the fascia of the vastus medialis.<sup>16 20 21</sup> However, the spread of dye injectate in cadavers may not be replicated in living subjects because of differences in tissue elasticity.<sup>16 17</sup>

Some secondary outcomes from our study differed from previous studies.<sup>4 5 9-11</sup> Compared with a continuous femoral triangle block, continuous proximal and distal ACBs reduced rescue analgesic requirements during the first 24 hours. Analgesia on POD2 was improved with continuous distal ACB compared with continuous femoral triangle block and continuous proximal ACB. Theoretically, continuous administration of local anesthetics to the distal AC could spread into the popliteal fossa and contribute to analgesic effects on posterior knee pain. The addition of supplementary blocks targeting posterior knee pain to ACB has been shown to improve analgesic outcomes after TKA.<sup>22-24</sup> Most previous ACB studies regarding have used periarticular local anesthetic infiltration.<sup>4 5 9 10</sup> Among them, one study compared a distal ACB with a proximal location, but unlike ours, they investigated clinical outcomes up to POD1.<sup>5</sup> At our institute, periarticular local anesthetic infiltration is a component of multimodal joint pathways and the posterior knee capsule was among the tissues routinely infiltrated. The analgesic benefit observed on POD2 may be attributed to the continuous distal ACB blocking posterior knee pain after the effect of periarticular local anesthetic infiltration disappeared.

There were no differences in quadriceps strength among groups, similar to previous studies of continuous femoral triangle blocks and ACBs.<sup>4 5 9–11</sup> We found a significant decrease in quadriceps strength compared with preoperative baseline values in all

groups. Postoperative quadriceps muscle strength can be affected by acute postoperative pain and surgical factors. This study was not powered to detect differences in this secondary outcome.

This study has several limitations. First, continuous catheter blocks were supplemented with periarticular local anesthetic infiltration, which might mask the analgesic benefit of distal ACB due to local anesthetic coverage for the posterior knee capsule on POD1. Second, the results apply to the specific local anesthetic type, basal rate, bolus volume, and lockout time used in this study. Third, more time was required to insert the catheter into the distal AC using a short-axis lateral to medial approach to avoid nerve damage by accurately tracing the saphenous nerve, which was not readily visible at this level. Finally, we determined each catheter insertion location in relation to individual femur length based on both ultrasound and radiographic images. However, whether the femur length ratio may be useful in determining the catheter insertion site in various races warrants further research.

In conclusion, continuous femoral triangle, proximal AC, and distal AC blocks provide comparable postoperative analgesia after TKA when periarticular local anesthetic infiltration is routinely administered. Quadriceps strength and opioid consumption did not differ among three continuous catheter blocks.

**Acknowledgements** The authors would like to thank Medical Illustration & Design, a part of the Medical Research Support Services of Yonsei University College of Medicine, for all the artistic support related to this work. The authors also would like to thank the biostatisticians employed for their statistical comments and analysis.

**Contributors** BL and YSC contributed to study conception and design. BL, SJP, KKP, HJK, YSL, and YSC contributed to study conduct. BL and SJP contributed to data analysis. BL, SJP, KKP, and YSC contributed to manuscript preparation. YSC is a guarantor.

# **Original research**

**Funding** This work was supported by a faculty research grant from the Yonsei University College of Medicine (grant number: 6-2020-0075).

Disclaimer The funder was not involved in the study.

Competing interests None declared.

#### Patient consent for publication Not applicable.

**Ethics approval** This study involves human participants and was approved by the Severance Hospital institutional review board on December 11, 2019 (protocol number: 4-2019-1038). This study was conducted in accordance with the principles of the Declaration of Helsinki. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

#### REFERENCES

- Grosu I, Lavand'homme P, Thienpont E. Pain after knee arthroplasty: an unresolved issue. Knee Surg Sports Traumatol Arthrosc 2014;22:1744–58.
- 2 Jiang H-H, Jian X-F, Shangguan Y-F, et al. Effects of enhanced recovery after surgery in total knee arthroplasty for patients older than 65 years. Orthop Surg 2019;11:229–35.
- 3 Memtsoudis SG, Cozowicz C, Bekeris J, et al. Peripheral nerve block anesthesia/ analgesia for patients undergoing primary hip and knee arthroplasty: recommendations from the International consensus on Anesthesia-Related outcomes after surgery (ICAROS) group based on a systematic review and meta-analysis of current literature. Reg Anesth Pain Med 2021;46:971–85.
- 4 Mariano ER, Kim TE, Wagner MJ, et al. A randomized comparison of proximal and distal ultrasound-guided adductor canal catheter insertion sites for knee arthroplasty. J Ultrasound Med 2014;33:1653–62.
- 5 Sztain JF, Khatibi B, Monahan AM, et al. Proximal versus distal continuous adductor canal blocks: does varying perineural catheter location influence analgesia? A randomized, Subject-Masked, controlled clinical trial. Anesth Analg 2018;127:240–6.
- 6 Bendtsen TF, Moriggl B, Chan V, et al. The optimal analgesic block for total knee arthroplasty. *Reg Anesth Pain Med* 2016;41:711–9.
- 7 Burckett-St Laurant D, Peng P, Girón Arango L, et al. The nerves of the adductor canal and the innervation of the knee: an anatomic study. *Reg Anesth Pain Med* 2016;41:321–7.
- 8 Anagnostopoulou S, Anagnostis G, Saranteas T, et al. Saphenous and infrapatellar nerves at the adductor canal: anatomy and implications in regional anesthesia. Orthopedics 2016;39:e259–62.
- 9 Chuan A, Lansdown A, Brick KL, et al. Adductor canal versus femoral triangle anatomical locations for continuous catheter analgesia after total knee arthroplasty: a multicentre randomised controlled study. Br J Anaesth 2019;123:360–7.

- 10 Meier AW, Auyong DB, Yuan SC, et al. Comparison of continuous proximal versus distal adductor canal blocks for total knee arthroplasty: a randomized, double-blind, Noninferiority trial. *Reg Anesth Pain Med* 2018;43:36–42.
- 11 Fei Y, Cui X, Chen S, et al. Continuous block at the proximal end of the adductor canal provides better analgesia compared to that at the middle of the canal after total knee arthroplasty: a randomized, double-blind, controlled trial. *BMC Anesthesiol* 2020;20:260.
- 12 Sabharwal S, Zhao C, McKeon JJ, et al. Computed radiographic measurement of limblength discrepancy. full-length standing anteroposterior radiograph compared with scanogram. J Bone Joint Surg Am 2006;88:2243–51.
- 13 Kim DH, Lin Y, Goytizolo EA, et al. Adductor canal block versus femoral nerve block for total knee arthroplasty: a prospective, randomized, controlled trial. Anesthesiology 2014;120:540–50.
- 14 Nielsen S, Degenhardt L, Hoban B, et al. A synthesis of oral morphine equivalents (OME) for opioid utilisation studies. Pharmacoepidemiol Drug Saf 2016;25:733–7.
- 15 Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- 16 Tran J, Chan VWS, Peng PWH, et al. Evaluation of the proximal adductor canal block injectate spread: a cadaveric study. Reg Anesth Pain Med 2020;45:124–30.
- 17 Jæger P, Jenstrup MT, Lund J, et al. Optimal volume of local anaesthetic for adductor canal block: using the continual reassessment method to estimate ED95. Br J Anaesth 2015;115:920–6.
- 18 Elazab EEB. Morphological study and relations of the fascia vasto-adductoria. Surg Radiol Anat 2017;39:1085–95.
- 19 Andersen HL, Andersen SL, Tranum-Jensen J. The spread of injectate during saphenous nerve block at the adductor canal: a cadaver study. *Acta Anaesthesiol Scand* 2015;59:238–45.
- 20 Johnston DF, Black ND, Cowden R, et al. Spread of dye injectate in the distal femoral triangle versus the distal adductor canal: a cadaveric study. *Reg Anesth Pain Med* 2019;44:39–45.
- 21 Runge C, Moriggl B, Børglum J, et al. The spread of ultrasound-guided Injectate from the adductor canal to the Genicular branch of the posterior obturator nerve and the popliteal plexus: a cadaveric study. *Reg Anesth Pain Med* 2017;42:725–30.
- 22 Ochroch J, Qi V, Badiola I, et al. Analgesic efficacy of adding the IPACK block to a multimodal analgesia protocol for primary total knee arthroplasty. *Reg Anesth Pain Med* 2020;45:799–804.
- 23 Kim DH, Beathe JC, Lin Y, et al. Addition of infiltration between the popliteal artery and the capsule of the posterior knee and adductor canal block to Periarticular injection enhances postoperative pain control in total knee arthroplasty: a randomized controlled trial. Anesth Analg 2019;129:526–35.
- 24 Hussain N, Brull R, Sheehy B, *et al.* Does the addition of iPACK to adductor canal block in the presence or absence of periarticular local anesthetic infiltration improve analgesic and functional outcomes following total knee arthroplasty? A systematic review and meta-analysis. *Reg Anesth Pain Med* 2021;46:713–21.