

# Correction: *Ultrasound-guided transverse abdominis plane block, ilioinguinal/iliohypogastric nerve block, and quadratus lumborum block for elective open inguinal hernia repair in children: a randomized controlled trial*

Priyadarshini K, Behera BK, Tripathy BB, *et al.* Ultrasound-guided transverse abdominis plane block, ilioinguinal/iliohypogastric nerve block, and quadratus lumborum block for elective open inguinal hernia repair in children: a randomized controlled trial. *Reg Anesth Pain Med* 2022;47:217–21.

On page 3, the first sentence in the first paragraph should read:

The median postoperative paracetamol consumption was 300 (157.5), 240 (127.5), and 247.5 (127.5) mg in the TAP, II/IH, and QLB groups, respectively.

One page 3, [table 1](#) should read:

**Table 1** Patient characteristics and block performance data

	Group TAP	Group II/IH	Group QLB
Age (year)	4.2 (5.3)	4.5 (4.9)	3.5 (4.5)
Weight (kg)	19 (6)	16 (8.5)	16.5 (8.5)

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.

*Reg Anesth Pain Med* 2022;47:e2. doi:10.1136/rapm-2021-103201corr1

