

Supplementary Table 2. Risk-adjusted odds of outcome measures for patients undergoing THA according to hospital safety net burden category (in quartiles).

<i>Safety net burden (quartiles)</i>	In-hospital mortality	Any complication	Cardiac complication	Pulmonary complication	Infectious complication	Intraoperative complication	LOS
First (lowest) quartile (reference)	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	(Reference)	(Reference)	(Reference)	(Reference)	(Reference)	(Reference)	(Reference)
Second quartile	1.34**	0.99	0.99	1.03	1.04	1.02	1.00
	(1.07, 1.67)	(0.89, 1.12)	(0.89, 1.12)	(0.89, 1.19)	(0.89, 1.22)	(0.79, 1.32)	(0.99, 1.00)
Third quartile	1.44**	1.03	1.02	1.05	1.05	1.06	1.03***
	(1.14, 1.82)	(0.92, 1.16)	(0.91, 1.15)	(0.91, 1.22)	(0.90, 1.23)	(0.81, 1.38)	(1.02, 1.03)
Fourth (top) quartile	1.61**	1.14	1.09	1.15	1.29**	1.30	1.08***
	(1.21, 2.15)	(1.00, 1.30)	(0.94, 1.26)	(0.98, 1.37)	(1.07, 1.54)	(0.97, 1.76)	(1.07, 1.09)
N	500110	500188	500188	500188	500188	500188	500188

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Percents may not sum to 100 due to rounding and missing values. All numbers reported are n (%), odds ratios (95% CI) unless noted otherwise.

^^Median (interquartile range)

^^^ Incidence rate ratio (IRR), 95% confidence interval (CI)

Models were clustered on hospital as a level-two covariate (random effect) and were adjusted for primary insurance payer, race/ethnicity, quartile of median income, sex, age, select Elixhauser comorbidity measures, year of procedure, quartile of hospital THA procedure volume; and hospital state