Rapid sequence spinal must be the new norm for an emergency LSCS in this Covid pandemic

To the Editor

I read with great interest the letter titled 'New indication for an old anesthetic technique: could we consider now rapid

sequence spinal anesthesia in a COVID-19 time?' by Varandas JS *et al.*¹

The advocacy of regional anesthesia as a whole has gained a large momentum in this pandemic. Regional anesthesia, which is not automatically contraindicated in these patients, plays a prominent role since it reduces high-risk interventions, potential staff contamination, patients' recovery time and maximize the management and efficiency of the operating room times.²

Regional anesthesia is promoted in obstetric practice for reasons of safety.¹ Regional anesthesia assumes even a greater role due to the benefits it brings to both mother and newborn.²

As already mentioned in the parent article—The rapid sequence spinal technique was first described in 2003 as a new approach to the provision of spinal anesthesia for the most urgent obstetric cases.³ The rapid sequence spinal technique is a no-touch technique, which means performing a central neuraxial block without palpating anatomic landmarks previously.³

Emergency lower segment cesarean section (LSCS) can be categorized based on Royal College of Obstetricians and Gynaecologists (RCOG) guidelines (2010) into four categories.⁴ Category I and category II indicate maternal or fetal compromise where, there should be an urgent delivery and the decision delivery interval should be within 30 and 45 min, respectively. In 2011, National Institute for Health and Care Excellence UK guidelines suggested a decision to delivery interval of 30 min for category I emergency LSCS and both 30 min and 75 min to category II Emergency LSCS.⁵

In a covid suspect or confirmed covid positive patient the team delivering the patient will probably be wearing the personal protective equipment (PPE) according to the parent institute recommended protocols. The process of donning the PPE and getting the operation theater (OT) ready even in a category I LSCS scenario is going to be time consuming and can take upto 30 min. At the same time 'there is no emergency in a pandemic' probably holds true to make sure that adequate and strict protocols of covid infection control are followed for the safety of the covid team. Hence in order to compensate to a small extent for the time taken to adequately don the PPE and get the OT prepared, the concept of a rapid sequence spinal should be the new norm in an emergency LSCS in this pandemic. It is important to note that reduction of decision delivery interval requires effective teamwork⁶ and cannot be over emphasized in this current situation.

Harshal D Wagh

Correspondence to Dr Harshal D Wagh, Anaesthesia, Kokilaben Dhirubhai Ambani Hospital, Mumbai, India; drhdw2701@gmail.com

Contributors HDW is the sole author.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial

purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

© American Society of Regional Anesthesia & Pain Medicine 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite Wagh HD. *Reg Anesth Pain Med* 2021;**46**:191–192.

Received 22 May 2020
Accepted 23 May 2020
Published Online First 2 June 2020

► <http://dx.doi.org/10.1136/rapm-2020-101572>

Reg Anesth Pain Med 2021;**46**:191–192.
doi:10.1136/rapm-2020-101725

ORCID iD

Harshal D Wagh <http://orcid.org/0000-0001-8538-8566>

REFERENCES

- 1 Varandas JS, Dias R, Mendes AB. New indication for an old anesthetic technique: could we consider now rapid sequence spinal anesthesia in a COVID-19 time? *Reg Anesth Pain Med* 2021;**46**:192.
- 2 Australian Society of Anaesthetists. *Anaesthesia and caring for patients during the COVID-19 outbreak*, 2020.
- 3 Kinsella SM, Girgih K, Scrutton MJL. Rapid sequence spinal anaesthesia for category-1 urgency caesarean section: a case series. *Anaesthesia* 2010;**65**:664–9.
- 4 Royal College of Obstetricians & Gynaecologists, the Royal College of Anaesthetists. Classification of urgency of caesarean section - a continuum of risk (Good Practice No. 11). Available: <https://www.rcog.org.uk/globalassets/documents/guidelines/goodpractice11classificationofurgency.pdf> [Accessed 19 May 2016].
- 5 National Institute for Health and Clinical Excellence. CG132 caesarean section. clinical guidelines for emergency Cs. Available: <http://www.guidance.nice.org.uk/> [Accessed 23 Jul 2016].
- 6 Siassakos D, Hasafa Z, Sibanda T, *et al.* Retrospective cohort study of diagnosis-delivery interval with umbilical cord prolapse: the effect of team training. *BJOG* 2009;**116**:1089–96.