

Evaluation of language analysis to summarize the literature: a comparison to traditional meta-analysis in primary hip and knee surgery

Joshua Myszewski ,¹ Emily Klossowski,² Kristopher M Schroeder³

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rapm-2021-103261>).

¹School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, USA

²College of Letters and Science, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin, USA

³Anesthesiology, University of Wisconsin-Madison, Madison, Wisconsin, USA

Correspondence to

Joshua Myszewski, University of Wisconsin-Madison School of Medicine and Public Health, Madison, Wisconsin, USA; jmyszewski@wisc.edu

Received 15 October 2021
Accepted 22 November 2021
Published Online First
14 December 2021



► <http://dx.doi.org/10.1136/rapm-2021-103381>



© American Society of Regional Anesthesia & Pain Medicine 2022. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Myszewski J, Klossowski E, Schroeder KM. *Reg Anesth Pain Med* 2022;**47**:151–154.

ABSTRACT

Introduction Sentiment analysis, by evaluating written wording and its context, is a growing tool used in computer science that can determine the level of support expressed in a body of text using artificial intelligence methodologies. The application of sentiment analysis to biomedical literature is a growing field and offers the potential to rapidly and economically explore large amounts of published research and characterize treatment efficacy.

Methods We compared the results of sentiment analysis of 115 article abstracts analyzed in a recently published meta-analysis of peripheral nerve block usage in primary hip and knee arthroplasty to the conclusions drawn by the authors of the original meta-analysis.

Results A moderately positive outlook supporting the utilization of regional anesthesia for hip and knee arthroplasty was found in the 115 articles that were included for analysis, with 46% expressing positive sentiment, 35% expressing neutral sentiment, and 19% of abstracts expressing negative sentiment. This was well aligned with the conclusions reached by a previous meta-analysis of the same articles.

Discussion Sentiment analysis applied to the medical literature can rapidly evaluate large collections of published data and generate an impression of overall findings that are aligned with the findings of a traditional meta-analysis.

INTRODUCTION

As the volume of medical literature continues to grow, meta-analysis studies represent an increasingly used and important mechanism to distil the available published data and make conclusions based on larger patient sample sizes. The clinical utility of these analyses results is derived from strengthened supportive conclusions in those settings where study findings are congruent and in dismissal of findings that fail to be replicated across published studies. While meta-analyses represent a powerful mechanism to evaluate published study data, their conduct can be limited by the time required to locate and evaluate appropriate published studies, export data that are to be included, and perform a statistical analysis. As the number of questions and scenarios explored by clinicians expands, the time constraints associated with traditional meta-analysis studies may render these types of analyses unwieldy and impractical for practicing clinicians.

One potential option to address the growing size of literature on any topic is the utilization of artificial

intelligence as an alternative or complement to traditional systematic review and meta-analysis. More specifically, a method such as sentiment analysis or other natural language processing methods could be used to quickly assess the findings of large groups of published studies and serve as a mechanism to generate further hypothesis.

Sentiment analysis is a type of artificial intelligence that can classify a body of text based on the qualitative sentiment (ie, the tone expressed) expressed within it and output either a categorical sentiment score (ie, positive, negative, neutral) or a numerical score which is on a spectrum where -1 is very negative and 1 is very positive.¹ This type of analysis has been applied in a variety of fields in the past for assessing large-scale trends such as those in social media² or bodies of clinical trial literature.³ Furthermore, specific algorithms have been developed that facilitate sentiment analysis of clinical trial abstracts.⁴ This presents an intriguing avenue for quick assessment of the qualitative statements made by the authors of a study. However, sentiment analysis as an adjunct or complement to other methods of systemic analysis of biomedical literature has not yet been explored.

With this in mind, we performed a sentiment analysis of the clinical trial abstracts used in a recent meta-analysis evaluating peripheral nerve block anesthesia use in primary hip and knee arthroplasty. We then compared the results of this sentiment analysis to what was reported in the meta-analysis. The goal of this study was to compare the results of sentiment analysis with the more exhaustive process of systematic review and explore its application as an adjunct to such methods of literature review.

METHODS

In this study, the sentiment (ie, the tone or level of support) of articles cited in a recent systematic review of the literature evaluating the use of peripheral nerve block analgesia for primary hip and knee arthroplasty was determined using GAN-BioBERT sentiment analysis.^{4,5} These findings were then compared with those expressed in a recent meta-analysis evaluating analgesic techniques for total hip and knee arthroplasty.⁵ This algorithm is publicly available under open-source license.⁶

Data collection

The abstracts of published manuscripts included in the systematic review by Memtsoudis *et al* that had abstracts available in the PubMed database were

Table 1 Samples of phrases with each sentiment category and value as determined by GAN-BioBERT

Sample phrases	Sentiment classification
The results of this study were promising.	Positive
The results of this study showed that treatment was contraindicated.	Negative
There were no significant differences found between the treatment and control group.	Neutral

collected using the NCBI's Entrez E-Utilities API by identifying the associated PMID for each paper.⁵ Manuscripts that either did not have abstracts available or were not indexed in PubMed were excluded.

Subgroup analyses

From the abstracts, several subgroup analyses were also performed. The subgroups included division by anesthesia technique, study type, surgery type (ie, knee vs hip arthroplasty), risk of bias, and number of patients in the study.

For the bias-based subgroups, studies with a bias rating risk of 'high' for any study characteristic as determined by Memtsoudis *et al* using the Cochrane Risk of Bias Tool were included in the high risk of bias subgroup; all other studies were treated as having a low risk of bias.

The specific division for the subgroups divided by patients per study was studies with greater than the median number of patients of all the studies analyzed versus those that did not.

Sentiment analysis

The algorithm used for sentiment analysis in this study, GAN-BioBERT, was written based on a semisupervised version of the previously described bidirectional encoder representations from transformers (BERT) algorithm for natural language processing by Devlin *et al*.^{7,8} GAN-BioBERT was built by Myszewski *et al* specifically for categorically classifying the tone expressed in clinical abstracts as either positive, negative, or neutral. Examples of text classified into each of these categories are shown in table 1.

To understand the utility of GAN-BioBERT as it relates to this study, we first need to discuss its precursors, BERT and BioBERT. The process by which GAN-BioBERT comes from these previous methods is shown graphically in figure 1.

A detailed description of the original BERT algorithm is available elsewhere,⁷ but some discussion of the algorithm and

its derivations that led to the algorithm used in this study is important to understanding its applicability to clinical literature.

The original BERT algorithm is currently considered as one of the state-of-the-art methodologies for natural language processing with high levels of accuracy. This algorithm uses a methodology known as transfer learning wherein a pretrained language model for a particular domain (ie, biomedical literature) is first developed with an extremely large sample of text from the language domain being studied. The original BERT model by Devlin *et al* was trained on a set of 2.5 billion words of text from English Wikipedia as well as 800 million words of text from BooksCorpus, a large collection of English language novels.⁷ This original BERT model was designed to be suited for general language tasks, but not to understand the nuances and complicated language frequently used in biomedical literature.

This general model was then refined in a second 'fine-tuning' step with fewer samples for the task of interest, (ie, biomedical literature) in a method that is dependent on the appropriateness of the original general language model.

To make this general language model more applicable to biomedical literature BioBERT was developed in 2020 by Lee *et al*.⁹ This was done by further training the original BERT model with an additional 4.5 billion words from PubMed abstracts as well as 13.5 billion words from PubMed Central full-text articles to create the language model known as BioBERT,⁹ which is uniquely designed/appropriate for the nuances of the writing style and terminology used in biomedical and academic literature

The algorithm/language model used in this study, GAN-BioBERT, was developed by Myszewski *et al* by further fine-tuning the biomedically oriented language model BioBERT for the specific task of classifying clinical study abstract sentiment.

The term fine-tuning is used to describe the process of refining a pretrained language model, that is, BioBERT, for a particular task such as sentiment classification of clinical trial abstracts by providing a smaller task specific set of examples to the algorithm. Following this fine-tuning step, the algorithm's performance is assessed and can then be applied for the proposed application (ie, sentiment classification of biomedical abstracts).

The sentiment classifications made by the GAN-BioBERT algorithm used for this study coincided with the determination of clinicians 91.3% of the time for classifying the sentiment in clinical trial abstracts as positive, negative, or neutral. This was determined by comparing the sentiment classifications made by the algorithm to the categorical classifications made by a set of clinicians on a set of sample abstracts that contained an equal




Figure 1 A graphical representation of the development of GAN-BioBERT as it relates to this study. BERT, bidirectional encoder representations from transformers.

amount of positive, negative, and neutral abstracts as determined by the clinicians.⁴

It is important to note that on an individual article level, GAN-BioBERT only categorizes abstracts as positive, negative, or neutral. This three-category classification scheme was chosen to provide adequate detail without sacrificing an adequate level of accuracy. More finely grained algorithms for sentiment classification are significantly limited by drops in accuracy as the granularity of the sentiment classifications is increased. For example, the original BERT algorithm was correct only 55.5% of the time when used for a sentiment classifier with five categories, as compared with being correct 93.1% of the time for a two-category benchmark dataset.⁷ Given this limitation, GAN-BioBERT is only suited to large samples of articles and not for use on individual articles.

RESULTS

Of the 122 study abstracts included in the original meta-analysis, 115 were included for sentiment analysis. One article was excluded due to not having an abstract but was still indexed in PubMed and six other articles were excluded due to not being indexed within the PubMed database. Each study, alongside its corresponding sentiment value determined by the algorithm is provided in online supplemental appendix A1.

Of the 115 articles included for analysis, it was determined that 56 (46.1%) had positive sentiment, 22 (19.1%) had negative sentiment, and 50 (34.8%) were neutral using sentiment analysis. The entire runtime for the analysis program, including data gathering, sentiment classification, and the subgroup analysis, was 8 min and 39 s on a desktop computer.

For comparison, the findings of the meta-analysis by Memtsoudis *et al* expressed a generally positive outlook toward the application of peripheral nerve block analgesia for total hip and knee arthroplasty.⁵ This sentiment was determined qualitatively based on the general recommendation made in that study that the use of peripheral nerve block analgesia is recommended for hip and knee arthroplasty as it leads to improved clinical outcomes. This generally positive outlook in the meta-analysis is aligned with the findings determined using sentiment analysis where the greatest proportion of studies had positive sentiment. It is also important to note that a large amount of the studies included expressed neutral sentiment, indicating a more moderate positive outlook.

The results for each of the subgroup analyses are shown in table 2. The median number of patients per study was found to be 80 patients.

DISCUSSION

In this study, a sentiment analysis of the sources used by a recent meta-analysis evaluating the clinical efficacy of regional anesthesia procedures for hip and knee arthroplasty reached a similar conclusion to that of the meta-analysis. In doing so, this current study accomplished two important goals that warrant further discussion. First, this study demonstrated that the GAN-BioBERT sentiment analysis approach yielded results concurrent with the findings of the Memtsoudis *et al* meta-analysis. Second, this study demonstrated that sentiment analysis represents an efficient alternative and/or complement to meta-analysis studies, with the total runtime of the analysis program requiring fewer than 10 min.

This study's (and the GAN-BioBERT algorithm's) major limitation is created by the three-class classification system and manifests as an inability to draw conclusions without an adequate sample size. This limitation restricts the use of the algorithm such that it cannot be used to draw conclusions on an individual level and is most appropriately used on a large aggregative scale to identify publication trends or as a precursor to more in-depth assessment of academic literature. With this in mind, the algorithm's application is limited to assessing larger trends such as topic-specific sentiment, as well as how this sentiment varies across time or specialty. Additional examples of possible applications fitting with this limitation include identifying trends related to publication biases as was shown in the subgroup analysis where the high bias and low bias subgroups had significantly different findings. Ideally, as the technology of sentiment analysis continues to advance, more granular classification schemes will achieve acceptable accuracy levels and the large sample size requirement will be ameliorated.

This study does possess several other limitations that should be considered prior to widespread application of sentiment scoring for guiding clinical decision-making. First, this specific algorithm only examines vocabulary located within abstracts of published studies. This limitation is imposed secondary to a significant loss of classification accuracy when the technology is used to classify sentiment for longer length bodies of text such as those in the body of manuscripts.¹ Therefore, text within the body of the manuscript that may recommend tempered

Table 2 Proportion of positive, negative, and neutral abstracts in each subgroup

Subgroup	Sample (n)*	Positive	Negative	Neutral
All included studies	115	53 (46.1%)	22 (19.1%)	50 (34.8%)
Both general and neuraxial anesthesia	26	12 (46.2%)	7 (26.9%)	7 (26.9%)
Only general anesthesia	24	12 (50%)	7 (29.2%)	5 (20.8%)
Only neuraxial anesthesia	49	22 (44.9%)	5 (10.2%)	22 (44.9%)
Hip arthroplasty	27	12 (44.4%)	5 (18.5%)	10 (37.1%)
Knee arthroplasty	92	41 (44.6%)	19 (20.6%)	32 (34.8%)
Studies with >80 patients	55	17 (30.9%)	17 (30.9%)	21 (38.2%)
Studies with <80 patients	60	36 (60%)	5 (8%)	19 (32%)
High risk of bias studies	37	23 (62.2%)	3 (8.1%)	11 (29.7%)
Low risk of bias studies	78	30 (38.5%)	19 (24.4%)	29 (37.1%)
Observational studies	40	15 (37.5%)	11 (27.5%)	14 (35%)
Randomized control trials	75	38 (50.7%)	11 (14.7%)	26 (34.6%)

*These subgroups are not all mutually exclusive and there may be overlap between groups.

enthusiasm or shades of optimism would not be subject to evaluation via the reported evaluation method. Second, this study only used sentiment analysis as a mechanism to validate a single meta-analysis study. It is possible that further examination of diverse meta-analysis studies may not result in similar findings or may find that different specialties are more apt to use positive, negative, or neutral terminology within their scientific writing. Third, the use of sentiment analysis in this context is unable to determine the impact of the studied intervention on a variety of clinical outcome domains. For example, the study by Memtsoudis *et al* demonstrated that the use of peripheral nerve blocks in the setting of knee and hip arthroplasty reduced the risk of cognitive dysfunction, respiratory failure, cardiac complications, and surgical site infections. However, in some cases, sentiment analysis may align closer to clinical decision-making where a clinician plans to administer a ‘better’ option and not necessarily one that offers an improvement in any distinct outcome domain.

Finally, performing a sentiment analysis may be beyond the technology limitations of an individual and therefore may not represent a feasible mechanism of literature review for all providers. Furthermore, while this study showed that sentiment analysis could accelerate the process of assessing literature, the application of sentiment analysis will still require the same careful consideration of the sources used for a particular research question or topic as is currently used in the process of systematic analyses and meta-analyses. However, it is important to consider that meta-analysis and expert opinion are imperfect and potentially subject to biases as rules are variably applied to which manuscripts are included in the final analysis or opinion is based on local experience and training.

Briefly, this study successfully used sentiment analysis as a rapid and efficacious mechanism to reach the same conclusion as a previously published meta-analysis study. With this in mind, sentiment analysis shows promise as a clinical literature evaluation tool that can be added to the repertoire of methods researchers and clinicians use to perform their work, verify study findings, and generate novel hypotheses.

Twitter Kristopher M Schroeder @KristopherSchr6

Contributors JM performed analysis, data gathering, manuscript writing, and project conception. EK performed manuscript editing and writing, analysis, and interpretation of the results. KMS provided supervision, guidance and institutional support, performed manuscript writing, and assisted with project conception.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study does not involve human participants.

Provenance and peer review Not commissioned; externally peer reviewed.

ORCID iD

Joshua Myszewski <http://orcid.org/0000-0002-3690-7056>

REFERENCES

- Choi Y, Lee H. Data properties and the performance of sentiment classification for electronic commerce applications. *Inf Syst Front* 2017;19:993–1012.
- Ceron A, Curini L, Iacus SM, *et al*. Every tweet counts? How sentiment analysis of social media can improve our knowledge of citizens’ political preferences with an application to Italy and France. *New Media & Society* 2014;16:340–58.
- Xu J, Zhang Y, Wu Y, *et al*. Citation sentiment analysis in clinical trial papers. *AMIA Annu Symp Proc* 2015;2015:1334–41.
- Myszewski JJ, Klossowski E, Meyer P. Validating GAN-BioBERT: a methodology for assessing reporting trends in clinical trials. *arXiv preprint arXiv* 2021:2106.00665.
- Memtsoudis SG, Cozowicz C, Bekeris J, *et al*. Peripheral nerve block anesthesia/analgesia for patients undergoing primary hip and knee arthroplasty: recommendations from the International consensus on Anesthesia-Related outcomes after surgery (ICAROS) group based on a systematic review and meta-analysis of current literature. *Reg Anesth Pain Med* 2021;46:971–85.
- Myszewski J. jmyszewski/GANBioBERTMetaAnalysis: initial release v1.0. *GitHub release* 2021.
- Devlin J, Chang MW, Lee K. Bert: Pre-training of deep bidirectional transformers for language understanding. *arXiv preprint arXiv* 2018:1810.04805.
- Croce D, Castellucci G, Basili R. GAN-BERT: generative adversarial learning for robust text classification with a bunch of labeled examples. Proceedings of the 58th Annual Meeting of the Association for Computational Linguistics, 2020:2114–9.
- Lee J, Yoon W, Kim S, *et al*. BioBERT: a pre-trained biomedical language representation model for biomedical text mining. *Bioinformatics* 2020;36:1234–40.

The majority of this data was retrieved from supplemental 1 provided by Memtsoudis et. al.,[1] with the addition of the sentiment scores from this study. As a reminder, a score of 1 is positive, -1 is negative, and 0 is neutral. n/a is provided for studies that were excluded from sentiment analysis due to either not having an abstract available on PubMed or for not being indexed in the PubMed database.

Randomized Control Trials

A: Allocation concealment B: Blinding of outcome assessors C: Blinding of participants and personnel D: Incomplete outcome data E: Other sources of bias F: Selective outcome reporting G: Sequence Generation

Study	Ref	Year	Study Type	n	A	B	C	D	E	F	G	Sentiment
Aksoy	2	2014	THA	70	Unclear	Low	High	Low	Low	Low	Low	1
Amundson	3	2017	TKA GA NA	157	Low	Low	Low	Low	Low	Low	Low	-1
Andersen	4	2012	TKA NA	40	Low	Low	Low	Low	Low	Low	Low	0
Angers	5	2019	TKA GA	90	Low	Low	Low	Low	Low	Low	Low	-1
Ashraf	6	2013	TKA NA	40	Low	Low	Low	Low	Low	Low	Unclear	0
Bali	7	2016	TKA GA	68	Low	Low	High	Low	Low	Low	Low	0
Baranovi	8	2011	TKA NA	71	Unclear	Unclear	Unclear	High	Low	Low	Low	1
Barrington	9	2005	TKA NA	108	Low	High	High	Low	Low	Low	Low	0
Beausang	10	2016	TKA NA	96	Unclear	High	High	Low	Low	Low	Unclear	0
Biswas	11	2018	TKA NA	130	Low	Low	Low	Unclear	Low	Low	Low	0
Bogoch	12	2002	THA TKA GA	115	Low	Low	Low	Low	Low	Low	Unclear	-1
Bron	13	2018	THA NA	162	Low	Low	Low	Low	Low	Low	Unclear	0
Campbell	14	2008	TKA NA	56	Unclear	Low	Unclear	High	Low	Low	Low	1
Chan	15	2013	TKA GA NA	135	Low	Low	High	Unclear	Low	Low	Low	1
Chan	16	2014	TKA	135	Unclear	Low	Unclear	Low	Low	Low	Unclear	0
Chaumeron	17	2013	TKA NA	59	Low	Low	Low	Low	Low	Low	Low	1
Chelly	18	2001	TKA GA NA	92	High	Unclear	High	Low	Low	Low	Unclear	1
Chen	19	2017	THA TKA GA	90	Low	Low	Unclear	Low	Low	Low	Low	n/a
Fahs	20	2018	THA GA	99	Low	Low	Unclear	Low	Low	Low	Low	1
Fan	21	2016	TKA GA	157	Low	Low	Low	Low	Low	Low	Low	-1
Fan	22	2017	TKA GA	65	Unclear	Unclear	Unclear	Low	Low	Low	Unclear	n/a
Fenten	23	2018	TKA NA	80	Low	Low	Low	Low	Low	Low	Low	1

Gasanova	24	2019	THA GA	60	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	1
Gmez	25	2017	TKA NA	574	Unclear	Unclear	Unclear	Low	Low	Low	Unclear	Unclear	1
Good	26	2007	TKA	42	Low	Low	Low	Low	Low	Low	Unclear	Unclear	1
Goytizolo	27	2016	THA NA	90	Low	Low	Low	Low	Unclear	Low	Low	Unclear	0
Goytizolo	28	2020	TKA NA	111	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	0
Grosso	29	2018	TKA NA	102	Low	Low	Low	Unclear	Low	Low	Low	Unclear	0
Hua	30	2017	THA GA	60	Unclear	Unclear	Unclear	High	Low	Low	Low	n/a	
Johnson	31	2017	THA GA NA	159	Low	Low	High	Low	Low	Low	Low	Unclear	0
Kadic	32	2009	TKA NA	53	Low	Low	Unclear	Low	Low	Low	Low	Unclear	-1
Kampitak	33	2018	TKA NA	57	Low	Low	Low	Low	Low	Low	Low	Unclear	1
Kardash	34	2007	TKA NA	40	Unclear	Low	Low	Low	Low	Low	Unclear	Unclear	-1
Kayupov	35	2018	TKA GA NA	91	Low	High	High	High	Unclear	Low	Low	Unclear	0
Kearns	36	2016	THA NA	108	Low	Low	Low	Low	Low	Low	Low	Unclear	-1
Kendrii	37	2017	THA GA NA	30	Unclear	Unclear	Unclear	Low	Low	Low	Unclear	n/a	
Kovalak	38	2015	TKA NA	60	Low	Unclear	Unclear	Low	Low	Low	Low	Unclear	1
Kratz	39	2015	THA GA	52	Low	Low	High	High	Low	Low	Low	Unclear	1
Kuchlik	40	2017	THA NA	56	Low	Low	Low	Low	Low	Low	Low	Unclear	0
Kulkarni	41	2019	TKA NA	100	Low	Low	Low	Low	Low	Low	Low	Unclear	-1
Lee	42	2011	TKA NA	78	Low	Low	Low	Low	Low	Low	Low	Unclear	0
Lee	43	2012	TKA GA	40	Unclear	Unclear	Unclear	Low	Low	Low	Unclear	Unclear	-1
Leung	44	2018	TKA NA	70	Low	Low	Low	Low	Low	Low	Low	Unclear	1
Li	45	2017	TKA	53	Low	Low	Low	Low	Low	Low	Low	Unclear	1
Long	46	2006	TKA NA	70	Unclear	Unclear	Unclear	High	Low	Unclear	Unclear	n/a	
Lu	47	2017	TKA GA NA	57	Unclear	Low	High	Low	Low	Low	Low	n/a	
Lueznier	48	2020	TKA	139	Low	Low	Low	Low	Low	Low	Low	Unclear	-1
Ivarez	49	2017	TKA NA	39	High	High	High	Low	Low	Low	Low	Unclear	1
Marino	50	2009	THA NA	150	Low	High	Unclear	Low	Low	Low	Low	Unclear	1
Mei	51	2017	THA GA	132	Low	Low	Unclear	Low	Low	Unclear	Low	Unclear	1
Moghtadaei	52	2014	TKA NA	36	Low	Low	Low	Unclear	Low	Low	Low	Unclear	1
Nader	53	2012	TKA NA	62	Low	Unclear	Unclear	Low	Low	Low	Low	Unclear	1

Ng	54	2001	TKA GA	48	Low	Low	Low	Low	Low	Low	Low	Low	-1
Ng	55	2012	TKA GA	32	Low	Low	Low	Low	Low	Low	Low	Low	0
Nishio	56	2014	THA GA NA	19	Low	Unclear	Unclear	Low	Low	Low	Low	Low	1
Niskanen	57	2005	TKA NA	50	Unclear	High	High	Low	Low	Low	Low	Unclear	1
Peng	58	2014	TKA GA	280	Low	Unclear	High	Low	Low	Low	Low	Low	1
Reinhardt	59	2014	TKA NA	94	Low	Low	Low	Low	Low	Low	Low	Low	0
Rizk	60	2017	TKA GA	75	Low	Unclear	Unclear	Low	Low	Low	Low	Low	n/a
Safa	61	2014	TKA NA	68	Low	Low	Low	Low	Low	Low	Low	Low	0
Sahin	62	2014	TKA NA	104	Low	Low	Low	Low	Low	Low	Low	Low	1
Saine	63	2018	TKA NA	60	Low	Low	Low	Low	Low	Low	Low	Low	0
Seet	64	2006	TKA NA	37	Low	Unclear	Unclear	Low	Low	Low	Low	Unclear	0
Siddiqui	65	2007	THA GA	34	Low	High	High	Low	Low	Low	Low	Low	1
Singelyn	66	1998	TKA GA NA	30	Low	Unclear	Unclear	Low	Low	Low	Low	Low	1
Sites	67	2004	TKA NA	40	Low	Unclear	Unclear	Low	Low	Low	Low	Low	1
Sogbein	68	2017	TKA NA	70	Low	Low	Low	Low	Low	Low	Low	Low	1
Spangehl	69	2015	TKA GA	160	Low	High	High	Low	Low	Low	Low	Low	1
Stathellis	70	2017	TKA GA	50	Low	Unclear	Unclear	Low	Low	Low	Low	Low	1
Stevens	71	2000	THA GA	60	Unclear	Unclear	Low	Low	Low	Low	Low	Unclear	1
Sundarathiti	72	2009	TKA NA	61	Unclear	Unclear	Unclear	Low	Low	Low	Low	Unclear	1
Thybo	73	2016	THA NA	100	Low	Low	Low	Unclear	Low	Low	Low	Low	0
Toftdahl	74	2007	TKA NA	77	Low	Unclear	Unclear	Low	Low	Low	Low	Low	0
Tong	75	2019	TKA NA	40	Low	Low	Low	Low	Low	Low	Low	Low	0
Twyman	76	1990	THA GA	20	Unclear	Unclear	Unclear	Low	High	Low	Low	Unclear	1
Wall	77	2017	TKA GA NA	257	Low	Low	Unclear	Unclear	Low	Low	Unclear	Low	1
Wang	78	2019	TKA	90	Low	Low	Low	Low	Low	Low	Low	Low	0
Widmer	79	2012	TKA GA	55	Low	Low	Low	Low	Low	Low	Low	Low	1
Wu	80	2014	TKA NA	79	Low	High	High	Unclear	Low	Low	Low	Low	1
Yamamoto	81	2019	THA NA	53	Low	Low	Low	Low	Low	Low	Low	Low	0
Zhou	82	2018	TKA GA	40	Low	Low	Low	Low	Low	Low	Low	Low	0
Zinkus	83	2017	TKA NA	54	Low	Low	Low	Low	Low	Low	Low	Low	1

Observational Studies

A: Failure to adequately control for confounding B: Failure to develop and apply appropriate eligibility criteria C: Flawed measurement of exposure or outcome D: Incomplete follow-up

Study	Year	Ref	Study Type	Study Technique	Patient	A	B	C	D	Sentiment
Akkaya	2014	84	Case-control	TKA NA	27	High	Low	Low	High	-1
Alsheik	2020	85	Retrospective cohort	TKA GA NA	80	Unclear	Low	Low	Low	1
Antoni	2014	86	Retrospective cohort	TKA GA	98	High	High	Low	Low	-1
Asakura	2011	87	Retrospective cohort	TKA GA	40	High	Low	Low	Low	1
Beaupre	2012	88	Prospective cohort	TKA GA NA	39	High	Low	Low	Low	0
Cien	2015	89	Retrospective cohort	TKA	122	Low	Low	Low	Low	0
Danninger	2014	90	Retrospective cohort	THA TKA GA NA	530089	Unclear	Low	Unclear	Low	-1
DeRuyter	2006	91	Prospective cohort	TKA GA NA	50	High	Low	Unclear	Low	1
Duncan	2013	92	Retrospective cohort	TKA GA NA	108	High	Unclear	Low	Low	1
Fetherston	2011	93	Prospective cohort	THA TKA	52	Unclear	Low	Low	Unclear	0
Fukuda	2020	94	Retrospective cohort	TKA GA	5094	Low	Low	Low	Low	-1
Green	2018	95	Retrospective cohort	THA GA	20	Unclear	Low	Low	Low	0
Gwam	2018	96	Retrospective cohort	TKA	110	Unclear	Low	Low	Low	-1
Henson	2019	97	Retrospective cohort	TKA	144	Unclear	Low	Low	Low	0
Horn	2015	98	Retrospective cohort	TKA	32	Unclear	Low	Low	High	1
Jacob	2011	99	Retrospective cohort	TKA GA NA	8590	Unclear	Low	Low	Low	-1
Jacob	2011a	100	Retrospective cohort	THA	9844	Unclear	Unclear	Low	Low	-1
Kim	2012	101	Retrospective cohort	TKA NA	80	High	Low	Low	Low	0
Kinjo	2012	102	Prospective cohort	TKA GA NA	81	Unclear	Low	Low	High	-1
Kirkness	2017	103	Retrospective cohort	TKA GA NA	268	Low	Low	Low	Low	n/a
Kukreja	2019	104	Retrospective cohort	THA NA	71	Low	Low	Low	Low	1

Liu	2015	105	Retrospective cohort	TKA NA	1768	Unclear	Low	Low	Low	0
Lovald	2015	106	Retrospective cohort	TKA GA NA	35642	High	Low	Unclear	Unclear	1
Mclsaac	2017	107	Retrospective cohort	TKA GA NA	178214	Low	Low	Low	Low	0
Memtsoudis	2016	107	Retrospective cohort	TKA GA NA	719426	Low	Low	Low	Low	-1
Memtsoudis	2016	108	Retrospective cohort	THA GA NA	342726	Low	Low	Low	Low	-1
Peters	2006	109	Retrospective cohort	THA TKA GA NA	100	High	High	Low	High	0
Pope	2015	110	Retrospective cohort	TKA GA NA	294	High	Low	Low	Low	0
Raimer	2007	111	Prospective cohort	TKA GA NA	42	Unclear	Low	Low	High	0
Rajeev	2016	112	Prospective cohort	TKA GA NA	114	High	High	Low	Low	1
Rames	2019	113	Retrospective cohort	TKA	693	Low	Low	Low	Low	0
Roberts	2019	114	Retrospective cohort	TKA	236	Low	Low	Low	Low	1
Schmidt	2009	115	Retrospective cohort	TKA GA NA	200	Unclear	Unclear	Low	Low	-1
Schwab	2019	116	Retrospective cohort	TKA	224	Unclear	Low	Low	Low	1
Simonsen	2011	117	Prospective cohort	TKA NA	67	Unclear	High	Low	Low	1
Singelyn	1999	118	Prospective cohort	THA GA NA	1274	High	Low	Low	Low	1
Sporer	2016	119	Retrospective cohort	TKA NA	597	Unclear	Low	Low	Low	0
Sugar	2011	120	Prospective cohort	TKA NA	28	Unclear	Low	Low	Low	1
Suthersan	2015	121	Prospective cohort	TKA GA NA	46	Unclear	Low	Low	Low	1
Tetsunaga	2016	122	Retrospective cohort	THA GA	62	High	High	Low	Low	0
Willett	2019	123	Retrospective cohort	TKA	151	High	Low	Low	Low	1

References

1. Memtsoudis SG, Cozowicz C, Bekeris J, et al. Peripheral Nerve block anesthesia/analgesia for patients undergoing primary hip and knee arthroplasty: recommendations from the International Consensus on Anesthesia-Related Outcomes after Surgery (ICAROS) group based on a systematic review and meta-analysis of current literature. *Reg Anesth Pain Med*. 2021.
2. Aksoy M, Dostbil A, Ince I, et al. Continuous spinal anaesthesia versus ultrasound-guided combined psoas compartment-sciatic nerve block for hip replacement surgery in elderly high-risk patients: a prospective randomised study. *BMC Anesthesiol* 2014; 14: 99
3. Amundson AW, Johnson RL, Abdel MP, et al. A Three-arm Randomized Clinical Trial Comparing Continuous Femoral Plus Single-injection Sciatic Peripheral Nerve Blocks versus Periarticular Injection with Ropivacaine or Liposomal Bupivacaine for Patients Undergoing Total Knee Arthroplasty. *Anesthesiology* 2017; 126: 1139-50
4. Andersen HL, Gyrn J, Moller L, Christensen B, Zaric D. Continuous saphenous nerve block as supplement to single-dose local infiltration analgesia for postoperative pain management after total knee arthroplasty. *Reg Anesth Pain Med* 2012; 37
5. Angers M, Belzile É L, Vachon J, Beauchamp-Chalifour P, Pelet S. Negative Influence of femoral nerve block on quadriceps strength recovery following total knee replacement: A prospective randomized trial. *Orthop Traumatol Surg Res* 2019; 105: 633-7
6. Ashraf A, Raut VV, Canty SJ, McLauchlan GJ. Pain control after primary total knee replacement. A prospective randomised controlled trial of local infiltration versus single shot femoral nerve block. *The Knee* 2013; 20: 324-7
7. Bali C, Ozmete O, Eker HE, Hersekli MA, Aribogan A. Postoperative analgesic efficacy of fascia iliaca block versus periarticular injection for total knee arthroplasty. *J Clin Anesth* 2016; 35: 404-10
8. Baranovic S, Maldini B, Milosevic M, Golubic R, Nikolic T. Peripheral regional analgesia with femoral catheter versus intravenous patient controlled analgesia after total knee arthroplasty: a prospective randomized study. *Coll Antropol* 2011; 35: 1209-14
9. Barrington MJ, Olive D, Low K, Scott DA, Brittain J, Choong P. Continuous femoral nerve blockade or epidural analgesia after total knee replacement: a prospective randomized controlled trial. *Anesth Analg* 2005; 101: 1824-9
10. Beausang DH, Pozek JP, Chen AF, et al. A Randomized Controlled Trial Comparing Adductor Canal Catheter and Intraarticular Catheter After Primary Total Knee Arthroplasty. *The Journal of arthroplasty* 2016; 31: 298-301
11. Biswas A, Perlas A, Ghosh M, et al. Relative Contributions of Adductor Canal Block and Intrathecal Morphine to Analgesia and Functional Recovery After Total Knee Arthroplasty: A Randomized Controlled Trial. *Reg Anesth Pain Med* 2018; 43: 154-60 11 Bogoch ER, Henke M, Mackenzie T, Olschewski E, Mahomed NN. Lumbar paravertebral nerve block in the management of pain after total hip and knee arthroplasty: a randomized controlled clinical trial. *The Journal of arthroplasty* 2002; 17: 398-401
12. Bron JL, Verhart J, Sierevelt IN, De Vries D, Kingma HJ, Rademakers MV. No effect of double nerve block of the lateral cutaneous nerve and subcostal nerves in total hip arthroplasty: A randomized controlled trial. *Acta Orthop* 2018; 89: 272-7

13. Campbell A, McCormick M, McKinlay K, Scott NB. Epidural vs. lumbar plexus infusions following total knee arthroplasty: randomized controlled trial. *Eur J Anaesthesiol* 2008; 25: 502-7
14. Chan EY, Fransen M, Sathappan S, Chua NH, Chan YH, Chua N. Comparing the analgesia effects of single-injection and continuous femoral nerve blocks with patient controlled analgesia after total knee arthroplasty. *The Journal of arthroplasty* 2013; 28: 608-13
15. Chan EY, Teo YH, Assam PN, Fransen M. Functional discharge readiness and mobility following total knee arthroplasty for osteoarthritis: a comparison of analgesic techniques. *Arthritis Care Res* 2014; 66: 1688-94
16. Chaumeron A, Audy D, Drolet P, Lavigne M, Vendittoli PA. Periarticular injection in knee arthroplasty improves quadriceps function knee. *Clin Orthop* 2013; 471: 2284-95
16. Chelly JE, Greger J, Gebhard R, et al. Continuous femoral blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *J Arthroplasty* 2001; 16: 436-45
17. Chen C, Li M, Wang K, et al. Protective effect of combined general and regional anesthesia on postoperative cognitive function in older arthroplasty patients. *Int J Clin Exp Med* 2017; 10: 15453-8
18. Fahs AM, Koueiter DM, Kurdziel MD, Huynh KA, Perry CR, Verner JJ. Psoas Compartment Block vs Periarticular Local Anesthetic Infiltration for Pain Management After Anterior Total Hip Arthroplasty: A Prospective, Randomized Study. *The Journal of arthroplasty* 2018
19. Fan L, Yu X, Zan P, Liu J, Ji T, Li G. Comparison of Local Infiltration Analgesia With Femoral Nerve Block for Total Knee Arthroplasty: A Prospective, Randomized Clinical Trial. *The Journal of arthroplasty* 2016; 31: 1361-5
20. Fan R, Zhao L, Hong H. Effect of inhalation anesthesia combined with nerve block on improving postoperative cognitive function in elderly orthopedic patients. *Biomedical Research (India)* 2017; 28: 4485-9
21. Fenten MGE, Bakker SMK, Scheffer GJ, Wymenga AB, Stienstra R, Heesterbeek PJC. Femoral nerve catheter vs local infiltration for analgesia in fast track total knee arthroplasty: short-term and long-term outcomes. *Br J Anaesth* 2018; 121: 850-8
22. Gasanova I, Alexander JC, Estrera K, et al. Ultrasound-guided suprainguinal fascia iliaca compartment block versus periarticular infiltration for pain management after total hip arthroplasty: a randomized controlled trial. *Reg Anesth Pain Med* 2019; 44: 206-11
23. Ortiz-Gómez JR, Perepérez-Candel M, Vázquez-Torres JM, et al. Postoperative analgesia for elective total knee arthroplasty under subarachnoid anesthesia with opioids: Comparison between epidural, femoral block and adductor canal block techniques (with and without perineural adjuvants). Aprospective, randomized, clinical trial. *Minerva Anesthesiol* 2017; 83: 50-8
24. Good RP, Snedden MH, Schieber FC, Polachek A. Effects of a preoperative femoral nerve block on pain management and rehabilitation after total knee arthroplasty. *American journal of orthopedics (Belle Mead, NJ)* 2007; 36: 554-7
25. Goytizolo EA, Stundner O, Rúa SH, et al. The Effect of Regional Analgesia on Vascular Tone in Hip Arthroplasty Patients. *HSS journal : the musculoskeletal journal of Hospital for Special Surgery* 2016; 12: 125-31
26. Goytizolo EA, Lin Y, Kim DH, et al. Addition of Adductor Canal Block to Periarticular Injection for Total Knee Replacement: A Randomized Trial. *J Bone Joint Surg Am* 2019; 101: 812-20

27. Grosso MJ, Murtaugh T, Lakra A, et al. Adductor Canal Block Compared with Periarticular Bupivacaine Injection for Total Knee Arthroplasty: A Prospective Randomized Trial. *J Bone Joint Surg Am* 2018; 100: 1141-6
28. Hua X, Hu Y, Chen D, Xiao Y, Luo L. Efficacy and safety of ultrasound-guided fascia iliaca compartment block using dexmedetomidine combined with ropivacaine in aged patients undergoing hip replacement. *Int J Clin Exp Med* 2017; 10: 16484-91
29. Johnson RL, Amundson AW, Abdel MP, et al. Continuous Posterior Lumbar Plexus Nerve Block Versus Periarticular Injection with Ropivacaine or Liposomal Bupivacaine for Total Hip Arthroplasty: A Three-Arm Randomized Clinical Trial. *The Journal of bone and joint surgery American volume* 2017; 99: 1836-45
30. Kadic L, Boonstra MC, MC DEWM, Lako SJ, J VANE, Driessen JJ. Continuous femoral nerve block after total knee arthroplasty? *Acta Anaesthesiol Scand* 2009; 53: 914-20
31. Kampitak W, Tanavalee A, Ngarmukos S, Amarase C, Songthamwat B, Boonshua A. Comparison of Adductor Canal Block Versus Local Infiltration Analgesia on Postoperative Pain and Functional Outcome after Total Knee Arthroplasty: A Randomized Controlled Trial. *Malaysian orthopaedic journal* 2018; 12: 7-14
32. Kardash K, Hickey D, Tessler MJ, Payne S, Zukor D, Velly AM. Obturator versus femoral nerve block for analgesia after total knee arthroplasty. *Anesth Analg* 2007; 105: 853-8
33. Kayupov E, Okroj K, Young AC, et al. Continuous Adductor Canal Blocks Provide Superior Ambulation and Pain Control Compared to Epidural Analgesia for Primary Knee Arthroplasty: A Randomized, Controlled Trial. *J Arthroplasty* 2018; 33: 1040-4.e1
34. Kearns R, Macfarlane A, Grant A, et al. A randomised, controlled, double blind, non-inferiority trial of ultrasound-guided fascia iliaca block vs. spinal morphine for analgesia after primary hip arthroplasty. *Anaesthesia* 2016; 71: 1431-40 36 Kendrišić M, Šurbatović M, Djordjević D, Trifunović B, Jevdjić J. Analgesic efficacy and safety of four different anesthesia/postoperative analgesia protocols in patients following total hip arthroplasty. *Vojnosanit Pregl* 2017; 74: 814-20
35. Kovalak E, Doğan AT, Üzümcügil O, et al. A comparison of continuous femoral nerve block and periarticular local infiltration analgesia in the management of early period pain developing after total knee arthroplasty. *Acta orthopaedica et traumatologica turcica* 2015; 49: 260-6
36. Kratz T, Dette F, Schmitt J, Wiesmann T, Wulf H, Zoremba M. Impact of regional femoral nerve block during general anesthesia for hip arthroplasty on blood pressure, heart rate and pain control: A randomized controlled study. *Technology and health care : official journal of the European Society for Engineering and Medicine* 2015; 28: 257-62
37. Kuchálik J, Magnuson A, Lundin A, Gupta A. Local infiltration analgesia or femoral nerve block for postoperative pain management in patients undergoing total hip arthroplasty. A randomized, double-blind study. *Scandinavian journal of pain* 2017; 16: 223-30
38. Kulkarni MM, Dadheech AN, Wakankar HM, Ganjewar NV, Hedgire SS, Pandit HG. Randomized Prospective Comparative Study of Adductor Canal Block vs Periarticular Infiltration on Early Functional Outcome After Unilateral Total Knee Arthroplasty. *J Arthroplasty* 2019; 34: 2360-4

39. Lee AR, Choi DH, Ko JS, et al. Effect of combined single-injection femoral nerve block and patient-controlled epidural analgesia in patients undergoing total knee replacement. *Yonsei Med J* 2011; 52: 145-50
40. Lee JJ, Choi SS, Lee MK, Lim BG, Hur W. Effect of continuous psoas compartment block and intravenous patient controlled analgesia on postoperative pain control after total knee arthroplasty. *Korean J Anesthesiol* 2012; 62: 47-51
41. Leung P, Dickerson DM, Denduluri SK, et al. Postoperative continuous adductor canal block for total knee arthroplasty improves pain and functional recovery: A randomized controlled clinical trial. *J Clin Anesth* 2018; 49: 46-52
42. Li D, Tan Z, Kang P, Shen B, Pei F. Effects of multi-site infiltration analgesia on pain management and early rehabilitation compared with femoral nerve or adductor canal block for patients undergoing total knee arthroplasty: a prospective randomized controlled trial. *Int Orthop* 2017; 41: 75-83
43. Long WT, Ward SR, Dorr LD, Raya J, Boutary M, Sirianni LE. Postoperative pain management following total knee arthroplasty: a randomized comparison of continuous epidural versus femoral nerve infusion. *The journal of knee surgery* 2006; 19: 137-43
44. Lu Y, Huang HM, Yan J, Jiang H. Comparison of postoperative femoral nerve block, epidural block and intravenous patient-controlled analgesia in pain control and postoperative rehabilitation after total knee arthroplasty. *Int J Clin Exp Med* 2017; 10: 6680-7
45. Lutzner J, Gehring R, Beyer F. Slightly better pain relief but more frequently motor blockade with combined nerve block analgesia compared to continuous intraarticular analgesia after total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc* 2020; 28: 1169-76
46. Alvarez NER, Ledesma RJG, Hamaji A, Hamaji MWM, Vieira JE. Continuous femoral nerve blockade and single-shot sciatic nerve block promotes better analgesia and lower bleeding for total knee arthroplasty compared to intrathecal morphine: a randomized trial. *BMC Anesthesiol* 2017; 17: 64
47. Marino J, Russo J, Kenny M, Herenstein R, Livote E, Chelly JE. Continuous lumbar plexus block for postoperative pain control after total hip arthroplasty: a randomized controlled trial. *JBJS* 2009; 91: 29-37
48. Mei B, Zha H, Lu X, et al. Peripheral Nerve Block as a Supplement to Light or Deep General Anesthesia in Elderly Patients Receiving Total Hip Arthroplasty: A Prospective Randomized Study. *The Clinical journal of pain* 2017; 33: 1053-9
49. Moghtadaei M, Farahini H, Faiz SH, Mokarami F, Safari S. Pain Management for Total Knee Arthroplasty: Single-Injection Femoral Nerve Block versus Local Infiltration Analgesia. *Iranian Red Crescent medical journal* 2014; 16: e13247
50. Nader A, Kendall MC, Wixson RL, Chung B, Polakow LM, McCarthy RJ. A randomized trial of epidural analgesia followed by continuous femoral analgesia compared with oral opioid analgesia on short- and long-term functional recovery after total knee replacement. *Pain medicine (Malden, Mass)* 2012; 13: 937-47
51. Ng HP, Cheong KF, Lim A, Lim J, Puhaindran ME. Intraoperative single-shot "3-in-1" femoral nerve block with ropivacaine 0.25%, ropivacaine 0.5% or bupivacaine 0.25% provides comparable 48-hr analgesia after unilateral total knee replacement. *Canadian journal of anaesthesia = Journal canadien d'anesthesie* 2001; 48: 1102-8

52. Ng FY, Ng JK, Chiu KY, Yan CH, Chan CW. Multimodal periarticular injection vs continuous femoral nerve block after total knee arthroplasty: a prospective, crossover, randomized clinical trial. *The Journal of arthroplasty* 2012; 27: 1234-8
53. Nishio S, Fukunishi S, Juichi M, et al. Comparison of continuous femoral nerve block, caudal epidural block, and intravenous patient-controlled analgesia in pain control after total hip arthroplasty: a prospective randomized study. *Orthopedic reviews* 2014; 6
54. Niskanen RO, Strandberg N. Bedside femoral block performed on the first postoperative day after unilateral total knee arthroplasty: a randomized study of 49 patients. *The journal of knee surgery* 2005; 18: 192-6
55. Peng L, Ren L, Qin P, et al. Continuous Femoral Nerve Block versus Intravenous Patient Controlled Analgesia for Knee Mobility and Long-Term Pain in Patients Receiving Total Knee Replacement: A Randomized Controlled Trial. *Evidence-based complementary and alternative medicine : eCAM* 2014; 2014: 569107
56. Reinhardt KR, Duggal S, Umunna BP, et al. Intraarticular analgesia versus epidural plus femoral nerve block after TKA: a randomized, doubleblind trial. *Clin Orthop* 2014; 472: 1400-8
57. Rizk H, Hosni Y, Abdeldayem S. Combined adductor canal and sciatic nerve block compared with local intraarticular infiltration analgesia for total knee arthroplasty: A prospective blinded randomized controlled study. *Current Orthopaedic Practice* 2017; 28: 179-83
58. Safa B, Gollish J, Haslam L, McCartney CJ. Comparing the effects of single shot sciatic nerve block versus posterior capsule local anesthetic infiltration on analgesia and functional outcome after total knee arthroplasty: a prospective, randomized, double-blinded, controlled trial. *The Journal of arthroplasty* 2014; 29: 1149-53
59. Sahin L, Korkmaz HF, Sahin M, Atalan G. Ultrasound-guided single-injection femoral nerve block provides effective analgesia after total knee arthroplasty up to 48 hours. *Agri : Agri (Algoloji) Dernegi'nin Yayin organidir = The journal of the Turkish Society of Algology* 2014; 26: 113-8
60. Rousseau-Saine N, Williams SR, Girard F, et al. The Effect of Adductor Canal Block on Knee Extensor Muscle Strength 6 Weeks After Total Knee Arthroplasty: A Randomized, Controlled Trial. *Anesth Analg* 2018; 126: 1019-27
61. Seet E, Leong WL, Yeo AS, Fook-Chong S. Effectiveness of 3-in-1 continuous femoral block of differing concentrations compared to patient controlled intravenous morphine for post total knee arthroplasty analgesia and knee rehabilitation. *Anaesth Intensive Care* 2006; 34: 25-30
62. Siddiqui ZI, Cepeda MS, Denman W, Schumann R, Carr DB. Continuous Lumbar Plexus Block Provides Improved Analgesia With Fewer Side Effects Compared With Systemic Opioids After Hip Arthroplasty: A Randomized Controlled Trial. *Reg Anesth Pain Med* 2007; 32: 393-8
63. Singelyn FJ, Deyaert M, Joris D, Penderville E, Gouverneur JM. Effects of intravenous patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous three-in-one block on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty. *Anesth Analg* 1998; 87: 88-92

64. Sites BD, Beach M, Gallagher JD, Jarrett RA, Sparks MB, Lundberg CJ. A single injection ultrasound-assisted femoral nerve block provides side effect-sparing analgesia when compared with intrathecal morphine in patients undergoing total knee arthroplasty. *Anesth Analg* 2004; 99: 1539-43
65. Sogbein OA, Sondekoppam RV, Bryant D, et al. Ultrasound-Guided Motor-Sparing Knee Blocks for Postoperative Analgesia Following Total Knee Arthroplasty: A Randomized Blinded Study. *The Journal of bone and joint surgery American volume* 2017; 99: 1274-81
66. Spangehl MJ, Clarke HD, Hentz JG, Misra L, Blocher JL, Seamans DP. The Chitranjan Ranawat Award: Periarticular injections and femoral & sciatic blocks provide similar pain relief after TKA: a randomized clinical trial. *Clin Orthop* 2015; 473: 45-53
67. Stathellis A, Fitz W, Schnurr C, et al. Periarticular injections with continuous perfusion of local anaesthetics provide better pain relief and better function compared to femoral and sciatic blocks after TKA: a randomized clinical trial. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA* 2017; 25: 2702-7
68. Stevens RD, Van Gessel E, Flory N, Fournier R, Gamulin Z. Lumbar plexus block reduces pain and blood loss associated with total hip arthroplasty. *Anesthesiology* 2000; 93: 115-21
69. Sundarathiti P, Ruananukul N, Channum T, et al. A comparison of continuous femoral nerve block (CFNB) and continuous epidural infusion (CEI) in postoperative analgesia and knee rehabilitation after total knee arthroplasty (TKA). *Journal of the Medical Association of Thailand = Chotmai het thangphaet* 2009; 92: 328-34
70. Thybo KH, Schmidt H, Hägi-Pedersen D. Effect of lateral femoral cutaneous nerve-block on pain after total hip arthroplasty: a randomised, blinded, placebo-controlled trial. *BMC anesthesiology* 2016; 16: 21
71. Toftdahl K, Nikolajsen L, Haraldsted V, Madsen F, Tønnesen EK, Søballe K. Comparison of peri- and intraarticular analgesia with femoral nerve block after total knee arthroplasty: a randomized clinical trial. *Acta Orthop* 2007; 78: 172-9
72. Tong QJ, Lim YC, Tham HM. Comparing adductor canal block with local infiltration analgesia in total knee arthroplasty: A prospective, blinded and randomized clinical trial. *J Clin Anesth* 2018; 46: 39-43
73. Twyman R, Kirwan T, Fennelly M. Blood loss reduced during hip arthroplasty by lumbar plexus block. *The Journal of bone and joint surgery British volume* 1990; 72: 770-1
74. Wall PDH, Parsons NR, Parsons H, et al. A pragmatic randomised controlled trial comparing the efficacy of a femoral nerve block and periarticular infiltration for early pain relief following total knee arthroplasty. *The bone & joint journal* 2017; 99-B: 904-11
75. Wang Q, Yue Y, Li D, Yang Z, Yeersheng R, Kang P. Efficacy of Single-Shot Adductor Canal Block Combined With Posterior Capsular Infiltration on Postoperative Pain and Functional Outcome After Total Knee Arthroplasty: A Prospective, Double-Blind, Randomized Controlled Study. *J Arthroplasty* 2019; 34: 1650-5
76. Widmer BJ, Scholes CJ, Pattullo GG, Oussedik SI, Parker DA, Coolican MR. Is femoral nerve block necessary during total knee arthroplasty?: a randomized controlled trial. *J Arthroplasty* 2012; 27: 1800-5

77. Wu JW, Wong YC. Elective unilateral total knee replacement using continuous femoral nerve blockade versus conventional patient-controlled analgesia: perioperative patient management based on a multidisciplinary pathway. *Hong Kong medical journal = Xianggang yi xue za zhi* 2014; 20: 45-51
78. Yamamoto N, Sakura S, Noda T, et al. Comparison of the postoperative analgesic efficacies of intravenous acetaminophen and fascia iliaca compartment block in hip fracture surgery: A randomised controlled trial. *Injury* 2019; 50: 1689-93
79. Zhou M, Ding H, Ke J. Adductor canal block in combination with posterior capsular infiltration on the pain control after TKA. *Ir J Med Sci* 2018; 187: 465-71
80. Zinkus J, Mockutė L, Gelmanas A, Tamošiūnas R, Vertelis A, Macas A. Comparison of 2 Analgesia Modalities in Total Knee Replacement Surgery: Is There an Effect on Knee Function Rehabilitation? *Medical science monitor : international medical journal of experimental and clinical research* 2017; 23: 3019-25
81. Akkaya A, Tekelioglu UY, Demirhan A, et al. Ultrasound-guided femoral and sciatic nerve blocks combined with sedoanalgesia versus spinal anesthesia in total knee arthroplasty. *Korean J Anesthesiol* 2014; 67: 90-5
82. Alsheikh KA, Alkhelaifi AS, Alharbi MK, et al. Adductor canal blockade versus continuous epidural analgesia after total knee joint replacement: A retrospective cohort study. *Saudi journal of anaesthesia* 2020; 14: 38-43
83. Antoni M, Jenny JY, Noll E. Postoperative pain control by intra-articular local anesthesia versus femoral nerve block following total knee arthroplasty: impact on discharge. *Orthopaedics & traumatology, surgery & research : OTSR* 2014; 100: 313-6
84. Asakura Y, Tsuchiya H, Mori H, Yano T, Kanayama Y, Takagi H. Reduction of the incidence of development of venous thromboembolism by ultrasound-guided femoral nerve block in total knee arthroplasty. *Korean journal of anesthesiology* 2011; 61: 382-7
85. Beaupre LA, Johnston DB, Dieleman S, Tsui B. Impact of a preemptive multimodal analgesia plus femoral nerve blockade protocol on rehabilitation, hospital length of stay, and postoperative analgesia after primary total knee arthroplasty: a controlled clinical pilot study. *ScientificWorldJournal* 2012; 2012: 273821
86. Cien AJ, Penny PC, Horn BJ, Popovich JM, Taunt CJ. Comparison Between Liposomal Bupivacaine and Femoral Nerve Block in Patients Undergoing Primary Total Knee Arthroplasty. *J Surg Orthop Adv* 2015; 24: 225-9
87. Danninger T, Rasul R, Poeran J, et al. Blood transfusions in total hip and knee arthroplasty: an analysis of outcomes. *ScientificWorldJournal* 2014; 2014: 623460
88. De Ruyter ML, Brueilly KE, Harrison BA, Greengrass RA, Putzke JD, Brodersen MP. A pilot study on continuous femoral perineural catheter for analgesia after total knee arthroplasty: the effect on physical rehabilitation and outcomes. *J Arthroplasty* 2006; 21: 1111-7
89. Duncan CM, Moeschler SM, Horlocker TT, Hanssen AD, Hebl JR. A self-paired comparison of perioperative outcomes before and after implementation of a clinical pathway in patients undergoing total knee arthroplasty. *Reg Anesth Pain Med* 2013; 38: 533-8
90. Fetherston CM, Ward S. Relationships between post operative pain management and short term functional mobility in total knee arthroplasty patients with a femoral nerve catheter: a preliminary study. *J Orthop Surg* 2011; 6: 7

91. Fukuda T, Imai S, Simoda S, Nakdera M, Horiguchi H. Comparison of peripheral nerve block with local infiltration analgesia regarding walking ability after total knee replacement: A retrospective, propensity-score matched-pair cohort study. *J Orthop Surg (Hong Kong)* 2020; 28: 2309499020931656
92. Stuart Green M, Ryan Hoffman C, Iqbal U, Olabisi Ives O, Hurd B. Transmuscular Quadratus Lumborum Block Reduces Length of Stay in Patients Receiving Total Hip Arthroplasty. *Anesth Pain Med* 2018; 8: e80233
93. Gwam CU, Mistry JB, Richards IV, et al. Does Addition of Adductor Canal Blockade to Multimodal Periarticular Analgesia Improve Discharge Status, Pain Levels, Opioid Use, and Length of Stay after Total Knee Arthroplasty? *The journal of knee surgery* 2018; 31: 184-8
94. Henson KS, Thomley JE, Lowrie LJ, Walker D. Comparison of Selected Outcomes Associated with Two Postoperative Analgesic Approaches in Patients Undergoing Total Knee Arthroplasty. *AANA J* 2019; 87: 51-7
95. Horn BJ, Cien A, Reeves NP, Pathak P, Taunt CJ. Femoral Nerve Block vs Periarticular Bupivacaine Liposome Injection After Primary Total Knee Arthroplasty: Effect on Patient Outcomes. *The Journal of the American Osteopathic Association* 2015; 115: 714-9
96. Jacob AK, Mantilla CB, Sviggum HP, Schroeder DR, Pagnano MW, Hebl JR. Perioperative nerve injury after total knee arthroplasty: regional anesthesia risk during a 20-year cohort study. *Anesthesiology* 2011; 114: 311-7
97. Jacob AK, Mantilla CB, Sviggum HP, Schroeder DR, Pagnano MW, Hebl JR. Perioperative nerve injury after total hip arthroplasty: regional anesthesia risk during a 20-year cohort study. *Anesthesiology* 2011; 115: 1172-8
98. Kim JH, Cho MR, Kim SO, Kim JE, Lee DK, Roh WS. A comparison of femoral/sciatic nerve block with lateral femoral cutaneous nerve block and combined spinal epidural anesthesia for total knee replacement arthroplasty. *Korean journal of anesthesiology* 2012; 62: 448-53
99. Kinjo S, Lim E, Sands LP, Bozic KJ, Leung JM. Does using a femoral nerve block for total knee replacement decrease postoperative delirium? *BMC Anesthesiol* 2012; 12: 4
100. Kirkness CS, Ren J, Asche CV. Significant improvement of mobility recovery in acute care patients after total knee arthroplasty with liposomal bupivacaine injectable suspension. *Journal of Acute Care Physical Therapy* 2017; 8: 11-9
101. Kukreja P, MacBeth L, Sturdivant A, et al. Anterior quadratus lumborum block analgesia for total hip arthroplasty: a randomized, controlled study. *Reg Anesth Pain Med* 2019
102. Liu Q, Chelly JE, Williams JP, Gold MS. Impact of peripheral nerve block with low dose local anesthetics on analgesia and functional outcomes following total knee arthroplasty: a retrospective study. *Pain Med* 2015; 16: 998-1006
103. Lovald ST, Ong KL, Lau EC, Joshi GP, Kurtz SM, Malkani AL. Readmission and Complications for Catheter and Injection Femoral Nerve Block Administration After Total Knee Arthroplasty in the Medicare Population. *The Journal of arthroplasty* 2015; 30: 2076-81
104. McIsaac DI, McCartney CJ, Walraven CV. Peripheral Nerve Blockade for Primary Total Knee Arthroplasty: A Population-based Cohort Study of Outcomes and Resource Utilization. *Anesthesiology* 2017; 126: 312-20

105. Memtsoudis SG, Poeran J, Cozowicz C, Zubizarreta N, Ozbek U, Mazumdar M. The impact of peripheral nerve blocks on perioperative outcome in hip and knee arthroplasty—a population-based study. *Pain* 2016; 157: 2341-9
106. Peters CL, Shirley B, Erickson J. The effect of a new multimodal perioperative anesthetic regimen on postoperative pain, side effects, rehabilitation, and length of hospital stay after total joint arthroplasty. *The Journal of arthroplasty* 2006; 21: 132-8
107. Pope D, El-Othmani MM, Manning BT, Sepula M, Markwell SJ, Saleh KJ. Impact of Age, Gender and Anesthesia Modality on Post-Operative Pain in Total Knee Arthroplasty Patients. *Iowa Orthop J* 2015; 35: 92-8
108. Raimer C, Priem K, Wiese AA, et al. Continuous psoas and sciatic block after knee arthroplasty: good effects compared to epidural analgesia or i.v. opioid analgesia: a prospective study of 63 patients. *Acta Orthop* 2007; 78: 193-200
109. Rajeev A, Tumia N, Karn K, Kashyap S, Mayne D. Postoperative pain relief and functional outcome following total knee arthroplasty - a prospective comparative audit of three analgesic regimes. *Acta Orthop Belg* 2016; 82: 265-70
110. Rames RD, Barrack TN, Barrack RL, Nunley RM. Effect of Adductor Canal Block on Acute Perioperative Pain and Function in Total Knee Arthroplasty. *J Arthroplasty* 2019; 34: S164-S7
111. Roberts C, Foster D, Shi GG, et al. A Collaborative Approach to Pain Control Reduces In-hospital Opioid Use and Improves Range of Motion following Total Knee Arthroplasty. *Cureus* 2019; 11: e4678
112. Schmidt NR, Donofrio JA, England DA, McDonald LB, Motyka CL, Mileto LA. Extended-release epidural morphine vs continuous peripheral nerve block for management of postoperative pain after orthopedic knee surgery: a retrospective study. *AANA J* 2009; 77: 349-54
113. Schwab PE, Yombi J, Lavand'homme P, Thienpont E. Comparison of local infiltration analgesia with single injection femoral nerve block in total knee arthroplasty. *Acta Orthop Belg* 2019; 85: 122-9
114. Simonsen OH, Gorst-Rasmussen A, Simonsen AB, Jorgensen MB, Rathleff MS, Lundbye-Christensen S. Blood reinfusion combined with femoral nerve block in total knee replacement for patients with increased risk of bleeding. *Journal of orthopaedic surgery (Hong Kong)* 2011; 19: 64-8
115. Singelyn FJ, Gouverneur JM. Postoperative analgesia after total hip arthroplasty: i.v. PCA with morphine, patient-controlled epidural analgesia, or continuous "3-in-1" block?: a prospective evaluation by our acute pain service in more than 1,300 patients. *J Clin Anesth* 1999; 11: 550-4
116. Sporer SM, Rogers T. Postoperative Pain Management After Primary Total Knee Arthroplasty: The Value of Liposomal Bupivacaine. *The Journal of arthroplasty* 2016; 31: 2603-7
117. Sugar SL, Hutson LR, Shannon P, Thomas LC, Nossaman BD. Comparison of extended-release epidural morphine with femoral nerve block to patient-controlled epidural analgesia for postoperative pain control of total knee arthroplasty: a case-controlled study. *The Ochsner journal* 2011; 11: 17-21

118. Suthersan M, Pit S, Gordon L, Loman M, Pezzutti B, Freihaut R. Local infiltration analgesia versus standard analgesia in total knee arthroplasty. *Journal of orthopaedic surgery (Hong Kong)* 2015; 23: 198-201
119. Tetsunaga T, Tetsunaga T, Fujiwara K, Endo H, Ozaki T. Combination Therapy with Continuous Three-in-One Femoral Nerve Block and Periarticular Multimodal Drug Infiltration after Total Hip Arthroplasty. *Pain Res Manag* 2016; 2016: 1425201
120. Willett A, Lew R, Wardhan R. Is continuous proximal adductor canal analgesia with a periarticular injection comparable to continuous epidural analgesia for postoperative pain after Total Knee Arthroplasty? A retrospective study. *Romanian journal of anaesthesia and intensive care* 2019; 26: 9-15