



Editor in Chief

Brian D Sites (USA)

Associate Editor-in-Chief for Statistics

Matthew A Davis (USA)

Executive Editors

- Richard Brull (Canada)
Chad M Brummett (USA)
Asokumar Buvanendran (USA)
Xavier Capdevila (France)
Steven P Cohen (USA)
Yun Guan (USA)
Ruth Landau (USA)
Stavros Memtsoudis (USA)
Anahi Perlas (Canada)
De Q H Tran (Canada)
Christopher L Wu (USA)

Editorial Office

BMJ Publishing Group Ltd, BMA House, Tavistock Square, London WC1H 9JR, UK
T: +1 919 650 1459
F: +44 (0)20 7383 6668
E: info.rapm@bmj.com

Impact factor: 5.564

Disclaimer: The Editor of Regional Anesthesia and Pain Medicine has been granted editorial freedom and Regional Anesthesia and Pain Medicine is published in accordance with editorial guidelines issued by the World Association of Medical Editors and the Committee on Publication Ethics.

Copyright: © 2022 American Society of Regional Anesthesia and Pain Medicine. All rights reserved; no part of this publication may be reproduced in any form without permission of Regional Anesthesia and Pain Medicine.

Regional Anesthesia and Pain Medicine is published by BMJ Publishing Group Ltd, typeset by Exeter Premedia Services Private Ltd, Chennai, India and printed in the UK on acid-free paper.

Regional Anesthesia & Pain Medicine ISSN 1472-0205 (USPS 580 990). BMA House, Tavistock Square, London WC1H 9JR, UK. The US annual subscription price is \$578.

Airfreight and mailing in the USA by agent named World Container Inc, 150-15, 183rd Street, Jamaica, NY 11413, USA. Periodicals postage paid at Brooklyn, NY 11256. US POSTMASTER: Send address changes to Regional Anesthesia & Pain Medicine, World Container Inc, 150-15, 183rd Street, Jamaica, NY 11413, USA.

Editorial

509 Looking backward to move forward in perioperative pain management?
E R Mariano, K El-Boghdadly

Special article

511 American Society of Regional Anesthesia and Pain Medicine contrast shortage position statement
L Kohan, Z Pellis, D A Provenzano, A C S Pearson, S Narouze, H T Benzon

Infographics

519 Effects of hypnosis versus enhanced standard of care on postoperative opioid use after total knee arthroplasty: the HYPNO-TKA randomised clinical trial—an infographic
R K Gupta, J Markovits, E S Schwenk

520 Retrospective cohort study of peripheral nerve blocks and general anesthesia versus general anesthesia alone: an infographic
E S Schwenk, R K Gupta, M Yoshimura

Regional anesthesia and acute pain

521 Comparing the effects of peripheral nerve block and general anesthesia with general anesthesia alone on postoperative delirium and complications in elderly patients: a retrospective cohort study using a nationwide database
M Yoshimura, H Shiramoto, M Koga, A Yoshimatsu, Y Morimoto

527 Comparison of postoperative analgesic effects of posterior quadratus lumborum block and intrathecal morphine in laparoscopic donor hepatectomy: a prospective randomized non-inferiority clinical trial
S Lee, R A Kang, G S Kim, M S Gwak, G-S Choi, J M Kim, J S Ko

534 Effects of hypnosis versus enhanced standard of care on postoperative opioid use after total knee arthroplasty: the HYPNO-TKA randomized clinical trial
J Markovits, O Blaha, E Zhao, D Spiegel

541 Comparison of lateral quadratus lumborum and lumbar plexus blocks for postoperative analgesia following total hip arthroplasty: a randomized clinical trial
T Kelly, C D Wolla, B J Wolf, E Hay, S Babb, S H Wilson

547 Effect of suprainguinal fascia iliaca compartment block on postoperative opioid consumption and functional recovery in posterolateral-approached total hip arthroplasty: a single-blind randomized controlled trial
M Carella, F Beck, N Piette, S Denys, W Kurth, J-P Lecoq, V L Bonhomme

554 Randomized comparison between perineural dexamethasone and combined perineural dexamethasone-dexmedetomidine for ultrasound-guided infraclavicular block
J Aliste, S Layera, D Bravo, G Aguilera, H Erpel, A Garcia, M Lizama, R J Finlayson, D Q Tran

Chronic and interventional pain

560 Tramadol prescribed at discharge is associated with lower odds of chronic opioid use after elective total joint arthroplasty
S Kiani, J Poeran, H Zhong, L A Wilson, L Poultsides, J Liu, S G Memtsoudis

Pediatric analgesia

566 Quantitative assessment of a pediatric nociception monitor in children under sevoflurane anesthesia
S Lebrun, J Boccara, E Cailliau, M Herbet, B Tavernier, I Constant, N Sabourdin

Research reports

571 Comparison of ultrasound-assistance versus traditional palpation method for placement of thoracic epidural catheters: a randomized controlled trial
C Arzola, M Ballei, Y Gleicher, A Malavade, Z Friedman

MORE CONTENTS ►



This article has been chosen by the Editor to be of special interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals open access scheme. See http://authors.bmj.com/open-access



Member since 2008

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org



When you have finished with this please recycle it

- 573** Trends in discontinuation of buprenorphine following elective orthopedic surgery: a national database analysis  
*A B Stone, H Zhong, J Poeran, J Liu, C Cozowicz, A Illescas, S G Memtsoudis*

---

### Brief technical report

- 576** Cryoanalgesia for shoulder pain: a motor-sparing approach to rotator cuff disease  
*A R Siogicza, P Peng*

---

### Case report

- 581** Percutaneous auricular nerve stimulation (neuromodulation) for the treatment of pain following outpatient surgery: a proof-of-concept case series  
*B M Ilfeld, J J Finneran IV, D Dalstrom, A M Wallace, B Abdullah, E T Said*



---

### Letters

- 586** Depth of cervical plexus block and diaphragmatic dysfunction  
*P Shao, C Han, Y Wang*
- 586** In reply to the letter to the editor by Wang *et al*  
*M Opperer, R Kaufmann, M Meissnitzer, F K Enzmann, C Dinges, W Hitzl, J Nawratil, A Koköfer*
- 587** Response to 'Efficacy of parasternal block to decrease intraoperative opioid use in coronary artery bypass surgery via sternotomy: a randomized controlled trial' by Bloc *et al*  
*S C Kim, S Anwar*
- 588** Parasternal block should be performed preoperatively, in response to Drs Seung Cheol Kim and Sibtain Anwar  
*S Bloc, C Quemeneur, F Le Sache, C Naudin*