

EP248

MEPIVACAINE DOSING FOR SPINAL ANESTHESIA IN PEDIATRIC ORTHOPEDIC SURGERY: A RETROSPECTIVE CHART REVIEW


Michelle Carley, Miriam Sheetz*, Justas Lauzadis, Haoyan Zhong, Kathryn DelPizzo. Department of Anesthesiology, Critical Care and Pain Management, Hospital for Special Surgery, New York, USA

10.1136/rapm-2023-ESRA.308


Background and Aims There is substantial literature on the use of spinal anesthesia in pediatric patients with bupivacaine, particularly in infants. Bupivacaine is a long-acting local anesthetic which is well suited to surgery in infants but less ideal for ambulatory surgery procedures in older children. Mepivacaine is an intermediate-acting agent commonly used for spinal anesthesia in adults and has potential benefits for use in older children. Currently, there are no published pediatric dosing guidelines for spinal mepivacaine. At Hospital for Special Surgery, mepivacaine is routinely used for spinal anesthesia in children. The aim of this study is to generate mepivacaine dosing guidelines based on milligrams per kilogram (mg/kg) and age.

Methods We performed a retrospective chart review of children who received mepivacaine for spinal anesthesia between 2016 to 2022.

Results The data extraction yielded 5,448 cases. Patient age ranged from 5 to 21 years. Mean surgery duration was 119 minutes (SD=48). Mean PACU length of stay was 222 minutes (SD=95). Weight in kilograms (kg) and mepivacaine dosage in milligrams (mg) was recorded for all patients (figure 1). The range and SD of total milligrams administered by age was also recorded (table 1). Median dosage in mg/kg of mepivacaine was calculated for each age group. Our analysis reveals that required dosage in mg/kg decreases by patient age and begins to plateau at age 15 (figure 2).



Abstract EP248 Figure 1 | Mean total mepivacaine dose administered by age



Abstract EP248 Figure 2 | Mepivacaine dosage in mg/kg reported as a function of patient age

Abstract EP248 Table 1 | Range of total milligrams of mepivacaine administered by age

Age (years)	Mean (mg)	SD	Dose range (mg)
5	37.5	5.3	30 – 45
6	42.5	5.9	37.5 – 52.5
7	44.8	4.3	37.5 – 54
8	42.5	2.7	37.5 – 45
9	46.8	5.9	33 – 60
10	48.8	6.5	33 – 63
11	50.6	7.2	30 – 66
12	51.8	6.8	37.5 – 66
13	53.2	6.9	37.5 – 69
14	53.5	7.1	37.5 – 75
15	54.6	6.8	30 – 75
16	54.6	7.3	30 – 75
17	54.7	7.5	30 – 75
18	54.6	7.9	30 – 75
19	54.9	7.2	30 – 75
20	55.3	7.0	30 – 75
21	55.9	6.6	37.5 – 75

Conclusions We describe mepivacaine dosage as a function of age and weight in children. As age and weight increase, a lower dose of mepivacaine per kg is required for spinal anesthesia.

IRB 2022-0706 approval

EP249

ABSTRACT WITHDRAWN

EP250

RECTUS SHEATH BLOCK ADDED TO PARASTERNAL BLOCK IMPROVES RESPIRATORY PERFORMANCE AFTER MEDIAN STERNOTOMY WITH DRAINAGE POSITIONING IN CARDIAC SURGERY PATIENTS


Alessandro Strumia*, Domenico Sarubbi, Annalaura Di Pumpo, Giuseppe Pascarella, Fabio Costa, Stefano Rizzo, Mariapia Stifano, Felice Eugenio Agrò. *Anaesthesia, Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy*

10.1136/rapm-2023-ESRA.310

Background and Aims Pain is usually severe after cardiac surgery and can limit respiratory function. Parasternal block is used to control this pain; anyway, the block effect is limited to the sternal region and do not cover upper abdominal quadrants, where pleural and mediastinal drainages are positioned. Rectus sheath block is an analgesic technique widely used in abdominal surgery.

Methods 5 patients underwent CABG through median sternotomy. With patients consent, we performed ultrasound guided bilateral parasternal block (ropivacaine 0,5% 40 ml + dexamethasone 2 mg) after induction and ultrasound guided bilateral rectus sheath block (ropivacaine 0,25% 20ml + dexamethasone 2mg) at the end of the surgery. Multimodal i. v. analgesia: ketorolac 90mg/24h and acetaminophen 1 gr 3/ die. Data regarded: perioperative pulmonary performance evaluated with the TriFlo Inspiratory Exerciser® and expressed in balls moved up during inspiration, pain during incentive spirometry at extubation/after 12 hours (0-10 NRS scale), opiates consumption.

Results Patients moved up a median of 2 (2-3) balls before surgery and a median of 2 (1-2) balls at extubation. 2 patients completely recovered respiratory function after 12 hours. Pain during spirometry at extubation was a median of 4 (3,5-5,5). Maximum pain in the first 12 hours was a median of 4 (3,5-5,5). Morphine consumption in the first 12 hours was a mean of 1 + 0,9 mg. No pulmonary complications occurred.



Abstract EP250 Figure 1 Rectus sheath block execution and local anaesthetic spread



Abstract EP250 Figure 2 The TriFlo Inspiratory Exerciser®

EP251

ANESTHETIC MANAGEMENT DURING LABOR AND SUBSEQUENT CESAREAN SECTION OF A PARTURIENT WITH DEVIC DISEASE (NEUROMYELITIS OPTICA): A CASE REPORT

Athanasia Tsaroucha, Christina Orfanou, Ailki Tympa-Grigoriadou, Thalys Asimakopoulos, Georgios Vaiopoulos, Aikaterini Melemini. *Department of Anesthesiology, School of Medicine, Aretaieion University Hospital, National and Kapodistrian University of Athens, Athens, Greece*

10.1136/rapm-2023-ESRA.311

Background and Aims Devic disease, or neuromyelitis optica, is a rare autoinflammatory demyelinating disease of the central nervous system, characterized by axonal damage, affecting mainly optic nerves and the spinal cord. The anesthetic management of a parturient suffering Devic disease in the delivery room, is presented.

Methods A 43-year-old, 90 kg, 167cm, G2P1 woman, diagnosed with Devic disease, presented for labor induction at 39 weeks of gestation. Initial neurologic symptoms, diplopia and facial nerve palsy, had developed during her first pregnancy and were diagnosed as brain stem syndrome in remission; the parturient received then uneventful epidural labor analgesia. A year later, Devic disease was diagnosed, further confirmed by