




## Results



**Abstract #33940 Figure 1** Perceived Lower Resistance to Injection using Pajunk® NRFit and Pajunk® Leur of the Sprotte® 24G x 90mm Spinal Needles (n=30) figure 2: Perceived Lower Resistance to Aspiration using Vygon® NRFit and Vygon® Leur of the Whitacre® 25G x 90mm Spinal Needles (n=30)



**Abstract #33940 Figure 1** Perceived Lower Resistance to Injection using Pajunk® NRFit and Pajunk® Leur of the Sprotte® 24G x 90mm Spinal Needles (n=30)



**Abstract #33940 Figure 2** Perceived Lower Resistance to Aspiration using Vygon® NRFit and Vygon® Leur of the Whitacre® 25G x 90mm Spinal Needles (n=30)

**Conclusions** Within the measure parameters, volunteers perceived a lower resistance to injection using the NRFit connectors. In contrast, they perceived lower resistance to aspiration using the leur connectors.

### #35832 AN ATYPICAL COMBINATION: SEDATION WITH DEXMEDETOMIDINE AND CONTINUOUS SPINAL ANESTHESIA FOR HIP FRACTURE ARTHROPLASTY IN A PATIENT WITH SEVERE DELIRIUM

<sup>1</sup>Rita Dinis, <sup>2</sup>Bárbara Sousa\*, <sup>1</sup>Ricardo Carvalho, <sup>3</sup>Andreia Puga. <sup>1</sup>Serviço de Anestesiologia, Hospital Prof. Doutor Fernando da Fonseca, Amadora, Portugal; <sup>2</sup>Serviço de Anestesiologia, Hospital Prof. Doutor Fernando da Fonseca, Sacavém, Portugal; <sup>3</sup>Serviço de Anestesiologia, Hospital Prof. Doutor Fernando da Fonseca, Lisboa, Portugal

10.1136/rapm-2023-ESRA.344

Please confirm that an ethics committee approval has been applied for or granted: Not relevant (see information at the bottom of this page)

**Application for ESRA Abstract Prizes:** I don't wish to apply for the ESRA Prizes

**Background and Aims** When the risks of approaching a difficult airway are high, regional anesthesia often becomes a wiser option. In order to avoid general anesthesia it becomes necessary to implement strategies that maximize the effectiveness of regional techniques, including optimizing patient cooperation and favoring regional techniques whose duration can be extended to meet the needs of prolonged surgery.

**Methods** Patient: 88-year-old female, with diabetes, hypertension, degenerative osteoarticular disease, dementia, and previous maxillectomy with ATM arthrodesis. Procedure: Hip arthroplasty due to hip fracture Anesthetic plan: A dexmedetomidine infusion was started 2 hours before coming to the operating room and was maintained throughout the procedure (0,1-0,4mcg/kg/h). A continuous spinal catheter (25G) was placed in L3/4, through an intrathecal Sprout needle (21G) – IntraLong (r) 21G/25G PAJUNK. An initial bolus of 2.5mg levobupivacaine 0,5% + 2.5mcg sufentanil was administered. The procedure lasted 2 hours and an additional dose of 1mg levobupivacaine 0,1% was given. The catheter was removed at the end of the procedure.

**Results** The patient remained in RASS -2, on spontaneous ventilation, without the need for additional oxygen supply and was hemodynamically stable throughout the procedure. The postoperative course was uneventful.

**Conclusions** In this case, avoiding the airway was highly desirable, but delirium could compromise patient cooperation during regional anesthesia. Sedation was necessary and dexmedetomidine was chosen because of its beneficial effect on delirium and respiratory stability. Continuous spinal anesthesia was chosen for its effectiveness in surgical anesthesia and due to the unpredictability of the duration of the procedure.

**Attachment** Consentimento informado sem dados do doente 2. pdf

### #36078 ASSESSING AND IMPROVING KNOWLEDGE OF EPIDURAL INFUSIONS AMONGST NON-ANAESTHETIC TRAINEES IN CRITICAL CARE

<sup>1</sup>Gillian Crowe\*, Ian Conrick-Martin. <sup>1</sup>Dublin, Ireland; <sup>2</sup>none, Dublin, Ireland

10.1136/rapm-2023-ESRA.345

Please confirm that an ethics committee approval has been applied for or granted: Not relevant (see information at the bottom of this page)

**Background and Aims** Epidural infusions provide good-quality analgesia after thoracic, abdominal and lower limb surgery and are commonly encountered in the post-operative patient in the high dependency unit. Management of epidural catheters and infusions are core skills for anaesthetic trainees, however recently the number of non-anaesthetic trainees working in critical care has increased. Exposure to epidural anaesthesia amongst this group is variable. Out of hours, in the absence of the pain team, the responsibility for management of the malfunctioning epidural may fall to a non- anaesthetic trainee. The aim of this project was to assess and improve knowledge of epidural infusions amongst non-anaesthetic trainees rotating through critical care.

**Methods** A 10-question multiple-choice questionnaire (MCQ) on epidural infusions was distributed amongst non-anaesthetic trainees rotating through critical care in our institution. An educational session was provided for a subset of this group after which they re-completed the MCQ. Pre- and post- education session scores were compared.

**Results** 15 non-anaesthetic trainees completed the MCQ, achieving an average score of 57% (range 40-70%). Eight trainees attended the education session. The mean post-education MCQ score improved to 95% (range 90-100%). The question most commonly answered incorrectly was related to calculating the length of catheter in the epidural space from the depth to loss of resistance and depth at skin. Key safety-related questions related to infusion rates, management of motor block and anti-coagulation were answered correctly by all participants following the education session.

**Conclusions** Familiarity with epidural infusions is mixed amongst non-anaesthetic trainees. A short educational session improves knowledge and familiarity amongst this group.

### #34490 TOTAL KNEE ARTHROPLASTY IN DOWN SYNDROME

About AlJabari\*. *Riyadh, Saudi Arabia*

10.1136/rapm-2023-ESRA.346

Please confirm that an ethics committee approval has been applied for or granted: Not relevant (see information at the bottom of this page)

**Application for ESRA Abstract Prizes:** I apply as an Anesthesiologist (Aged 35 years old or less)

**Background and Aims** We present a rare case of total knee arthroplasty in 32 year old down syndrome male patient, ASA 2, hypothyroidism presented with right knee pain scheduled for total knee arthroplasty under regional anesthesia.

**Methods** Pre anesthesia evaluation was assessed for airway management and best anesthesia plan according to his medical co-morbidities, blood results were within normal range. Cervical spine x-ray was requested. Thyroid function test revealed controlled treatment. His cardiac echo showed: The left ventricle is normal in size. There is normal left ventricular wall thickness. Left ventricular systolic function is normal. Ejection Fraction >55%. Left ventricular diastolic function is normal. The right ventricle is normal in size and function. Mildly

thickened Aortic valve leaflet. Mild to moderate aortic regurgitation. His electrocardiogram showed normal sinus rhythm. Our plan for his total knee arthroplasty is under spinal anesthesia along with ultrasound guided adductor canal catheter and IPACK infiltration for postoperative pain control. The duration of surgery lasted for two hours without any complications and 300 ml estimated blood loss and adequate urine output.

**Results** He was receiving multimodal analgesia of adductor canal ropivacaine injection daily through the catheter, paracetamol IV, morphine 2 mg IV PRN, NSAIDS once daily. He was discharged home without any complications and he is doing back his daily activities without any chronic knee pain.

**Conclusions** As far as we know, this is the first case presentation of total knee arthroplasty in Down syndrome as it is common for hip arthroplasty than knee.

### #36283 A RARE CASE OF PNEUMORRACHIS AFTER PLACEMENT OF AN EPIDURAL ELASTOMERIC DRUG INFUSION BALLOON (DIB)

<sup>1</sup>Paulo Correia\*, <sup>2</sup>Nelson Gomes, <sup>2</sup>Caroline Dahlem. <sup>1</sup>Anesthesiology, CHEDV, Santa Maria da Feira, Portugal; <sup>2</sup>Anesthesiology, CHEDV, Porto, Portugal

10.1136/rapm-2023-ESRA.347

Please confirm that an ethics committee approval has been applied for or granted: Not relevant (see information at the bottom of this page)

**Background and Aims** Pneumorrhachis is a rare complication of epidural analgesia and is most often asymptomatic. It can cause permanent deficit and differential diagnosis can be challenging, so clinicians should be aware of this entity.

**Results** A 74 year old woman was admitted to elective total knee replacement surgery. A L3/L4 spinal block using a paramedian approach was achieved after 2 attempts with a 25G quinke needle. An epidural catheter was placed with loss of resistance (LOR) to saline through L3/L4 intervertebral place, by single attempt. The procedure was uneventful and the sensitive and motor block reversed in the PACU. Before the transference to the ward, an epidural DIB was initiated with 0,1% ropivacaine, 5 mL/h. 4 hours later, the anesthesiologist was called for a bilateral sensitive and motor block up to T10 and urinary retention. After neurologist's assessment and DIB clamping, an MRI revealed intradural and extradural air collections, in locations compatible with the deficits presented. The patient was transferred to the hyperbaric medicine center with oxygen inhalation via a non-rebreather mask. Upon arrival, the deficits had completely reversed and it was decided to do 12 hours of normobaric oxygen therapy. Patient was discharged by 6th post-operative day and no other complications was observed.

**Conclusions** Pneumorrhachis after an epidural technique with LOR to saline is rare. Our most plausible hypothesis was that air could have been entrapped in the DIB. It usually gets reabsorbed spontaneously<sup>1</sup>. Nonoperative treatment includes hyperbaric oxygen therapy, which can lead to reabsorption of trapped air.