surgery under general anesthesia, were enrolled in a prospective, randomized, double-blind, single-centre, pilot study [NCT04530903].

Group. 1 received an ultrasound-pudendal block using 75µg of clonidine per side; group.2 0.5 ml of NaCl 0.9% per side; both groups received 10 ml of levobupivacaine 0,25%.

Primary outcome: total consumption of tramadol in PACU. Secondary outcomes: time between block and first analgesic demand; tramadol request within 24h; postoperative pain; satisfaction.

Continuous data were compared via T-test or Wilcoxon signed-rank test; means±st.deviation or medians are reported. For count data, Pearson Chi-Squared test was performed to compare proportions. $P < 0.05$ was considered significant.

Results Demographics and surgical data were comparable between groups. The groups were not statistically different for the primary outcome (group.1:100 mg; group.2:450 mg; $P = 0.1677$).

55.6% of group.2 requested tramadol in PACU ($P = 0.09677$) compared to 12.5% of group.1.

Within 24h, group.1 required more Tramadol ($P = 0.05$).

VAS longitudinal analysis showed a group-time effect with significantly higher pain within 24h in group.1 ($P = 0.0477$).

Satisfaction was not statically different between groups.

Conclusions Adding clonidine to levobupivacaine in pudendal block did not improve postoperative analgesia.

A large-scale study is needed to further support these results.

231

INFECTION AFTER KNEE ARTHROPLASTY, DO WE KNOW ALL RISK FACTORS?

B Sánchez Quirós*, LH Rocio, PH María Asunción. *Hospital Clínico Universitario de Valladolid, Valladolid, Spain*

10.1136/rapm-2021-ESRA.231

Background and Aims Non-steroidal anti-inflammatory drugs (NSAIDs) inhibit cyclooxygenase and it causes granulocytic function suppression (1). These effects could be found in relation to the increased incidence of infections and higher risk of bleeding (2).

Aim of this study was a descriptive analysis of risk factors for total knee arthroplasty, in order to evaluate the relationship between NSAIDs and perioperative infection (4).

Abstract 231 Table 1 Variables collected

Allergies
Personal history
Surgical history
Treatment with NSAIDs before and after the intervention.
Septic / aseptic replacement.
Joint liquid microbiology
Microbiology knee prosthesis
Antibiotic treatment
Administration of tranexamic acid.
Type of anesthesia
Analytical parameters: Hb, leukocytes, platelets, PCR.

Methods A descriptive observational study in 25 patients medical records who were scheduled for total knee arthroplasty in University Hospital of Valladolid. The following variables were collected (table 1). The study was approved by Hospital Clínico

Abstract 231 Table 2 Baseline clinical epidemiological characteristics of the sample

	n/N (%)
Female	17/25 (68%)
Diabetes Mellitus	7/25 (28%)
Hypertension	11/25 (44%)
Dyslipidemia	10/25 (40%)
Cardiopathy	10/25 (40%)
Obesity	4/25 (16%)
Periarticular infiltration	4/25 (16%)
Intraoperative administration of tranexamic acid	16/25 (64%)
Locoregional anesthesia	23/25 (92%)

Abstract 231 Table 3 Distribution of risk factors in both groups (septic/aseptic)

		ASEPTIC		SEPTIC	
		n	Percentage	n	Percentage
Gender	Men	7	50,0%	1	9,1%
	Female	7	50,0%	10	90,9%
Allergies	NO	11	78,6%	10	90,9%
	YES	3	21,4%	1	9,1%
Diabetes Mellitus	NO	10	71,4%	8	72,7%
	YES	4	28,6%	3	27,3%
Hypertension	NO	8	57,1%	6	54,5%
	YES	6	42,9%	5	45,5%
Dyslipidemia	NO	10	71,4%	7	63,6%
	YES	4	28,6%	4	36,4%
Cardiopathy	NO	8	57,1%	7	63,6%
	YES	6	42,9%	4	36,4%
Obesity	NO	12	85,7%	9	81,8%
	YES	2	14,3%	2	18,2%
Periarticular Infiltration	NO	12	85,7%	9	81,8%
	YES	2	14,3%	2	18,2%
Tranexamic acid	NO	3	21,4%	6	54,5%
	YES	11	78,6%	5	45,5%
Anesthesia	General	0	0,0%	2	18,2%
	Intradural	14	100,0%	9	81,8%

Universitario de Valladolid with number (CEIm PI 19/1438). All patients signed the inform consent to enter the study.

Results Statistically significant differences were found in the rate of infection with respect to sex; being more frequent the infections in the feminine sex ($p=0.042$). Marginally significant differences were found between the taking of NSAIDs and infection of the prosthesis ($p=0.1$) although there was no statistical significance between the taking of NSAIDs and the analytical parameters of infection. This study demonstrates that periarticular infiltration does not increase the risk of infection ($p < 0.05$)

Conclusions The use of NSAIDs has been associated with an immunosuppressive effect that leads to an increased risk of infection in total knee arthroplasty, so its use in the preoperative period should be limited. Female gender is other significant risk factor too. Our retrospective study has shown relationship between AINEs use and infection after knee arthroplasty. No increased risk of bleeding with AINEs in this sample.

232 CONTINUOUS ERECTOR SPINAE PLANE BLOCK FOR POSTOPERATIVE ANALGESIA AS AN ALTERNATIVE TO EPIDURAL CATHETER IN THORACIC SURGERY FOR PATIENTS WITH EMPYEMA: A CASE SERIES

C Simon*, N Parisi, G Samouri. *Clinique Saint-Pierre, Ottignies, Belgium*

10.1136/rapm-2021-ESRA.232

Background and Aims We present a case series of five patients undergoing video-assisted thoracoscopic surgery for empyema. In this infectious state, we chose to perform a continuous ESP because of the relative contraindication of epidural catheter. We performed it to prevent and relieve the thoracic pain but also to limit the opioids consumption and their side effects in our frailty patients.

Methods The plan was general anaesthesia combined with a continuous unilateral ESP block. Postoperative analgesia was achieved by continuous infusion + bolus of L-Bupivacaine 0.125%. Complementary analgesia was done with paracetamol and when possible NSAID. Patients received iv doses of piritramide in the recovery room, but no patient-controlled analgesia or other intravenous opioids was administered subsequently.

Results Most of the patients maintained a Numeric pain Rating Scale (NRS) between 0 and 3.

We removed the catheters the same day than the pleural drains, usually at day 4. In the postoperative period, no side

effects (motor blockade, urinary retention or hypotension) were noticed and patients attested high satisfaction scores. We emphasize the fact that none of them requested additional opioids.

Conclusions In cases of empyema, the ESP catheter combined with multimodal analgesia seems to be a safe and effective alternative to epidural analgesia for VATS surgery. This technique, easy to perform, allows an early mobilization of the patient regarding the low pain scores and the absence of side effects. Prospective randomized trials are needed to confirm the benefits of the ESPB compared to epidural or paravertebral bloc.

233 PECS II BLOCK FOR ONCOLOGIC BREAST SURGERY DURING PREGNANCY: A PATIENT-CENTERED CASE REPORT

¹J Pinho*, ²J Moita, ¹N Veiga. ¹Francisco Gentil Portuguese Oncology Institute of Lisbon, Department of Anesthesiology, Lisbon, Portugal; ²Setúbal Hospital Center, Department of Anesthesiology, Setúbal, Portugal

10.1136/rapm-2021-ESRA.233

Background and Aims Anesthesia and analgesia for surgery during pregnancy has focused on avoiding teratogenic drugs, preserving intrauterine homeostasis, and decreasing the risk of preterm labor. Data on the use of regional techniques and on patient preferences are still scarce. Based on a clinical case, we aimed to explore the patient experience of analgesia for oncologic breast surgery during pregnancy.

Methods A 19-weeks-pregnant 40-year-old woman with a diagnosis of breast cancer presented for mastectomy and sentinel lymph node biopsy. After discussion with the patient, an analgesia plan of post-induction pectoral type-2 block, acetaminophen, low-dose morphine, and postoperative acetaminophen, tramadol and *prn* metamizole was chosen. Acetaminophen and *prn* metamizole were prescribed home. Patient experience was elicited during the perioperative care and in a semi-structured interview at four-months.

Results On the first postoperative day, the patient reported only mild pain on movement and declined analgesia except for one administration of acetaminophen. After discharge, she reported significant pain on the surgical drain site. Until drain removal on the fourth postoperative day, she took acetaminophen and one *prn* metamizole. The main themes identified in the patient experience were a prioritization of fetal wellbeing, a concern for the effect of analgesic drugs on the fetus, and apprehension regarding nausea and sedation.

Conclusions Outcomes commonly measured in medical research are ultimately surrogates for outcomes that are more relevant to patients. This case suggests that a decreased need

DAILY EVOLUTION OF NRS

Abstract 232 Figure 1

Abstract 233 Table 1

Central questions	Main themes identified
<ul style="list-style-type: none"> • "How did you expect postoperative pain and analgesia would be?" • "How was your experience of postoperative pain and analgesia?" • "How would your ideal experience have been?" • "What should our goals be in the treatment of postoperative pain in pregnant women?" • "How should these goals be prioritized?" 	<ul style="list-style-type: none"> • Prioritization of fetal wellbeing over analgesia • Concern for the impact of analgesic drugs on the fetus • Valorization of the absence of nausea and sedation