



OPEN ACCESS

Recommendations for effective documentation in regional anesthesia: an expert panel Delphi consensus project

Hassan M Ahmed ¹, Benjamin P Atterton,² Gillian G Crowe,² Jaime L Barratta,³ Mark Johnson ⁴, Eugene Viscusi ³, Sanjib Adhikary ⁵, Eric Albrecht ^{6,7}, Karen Boretsky,⁸ Jan Boublik,⁹ Dara S Breslin,¹⁰ Kelly Byrne ¹¹, Alan Ch'ng,⁴ Alwin Chuan ¹², Patrick Conroy,¹³ Craig Daniel,¹⁴ Andrzej Daszkiewicz,¹⁵ Alain Delbos,¹⁶ Dan Sebastian Dirzu ¹⁷, Dmytro Dmytriiev ¹⁸, Paul Fennessy,¹⁹ H Barrie J Fischer,²⁰ Henry Frizelle,²¹ Jeff Gadsden,²² Philippe Gautier,²³ Rajnish K Gupta ²⁴, Yavuz Gürkan,²⁵ Harold David Hardman,²⁶ William Harrop-Griffiths,²⁷ Peter Hebbard,^{28,29} Nadia Hernandez,³⁰ Jakub Hlasny,³¹ Gabriella Iohom,^{32,33} Vivian H Y Ip ³⁴, Christina L Jeng ³⁵, Rebecca L Johnson ³⁶, Hari Kalagara,³⁷ Brian Kinirons,³⁸ Andrew Kenneth Lansdown,³⁹ Jody C Leng,^{40,41} Yean Chin Lim ⁴², Clara Lobo,⁴³ Danielle B Ludwin,⁴⁴ Alan James Robert Macfarlane ^{45,46}, Anthony T Machi ⁴⁷, Padraig Mahon,³² Stephen Mannion,⁴⁸ David H McLeod,⁴⁹ Peter Merjavy ⁵⁰, Aleksejs Miscuks,⁵¹ Christopher H Mitchell,⁵² Eleni Moka,⁵³ Peter Moran,³⁸ Ann Ngui,⁴ Olga C Nin,⁵⁴ Brian D O'Donnell,³² Amit Pawa ^{55,56}, Anahi Perlas ^{57,58}, Steven Porter ⁵⁹, John-Paul Pozek,⁶⁰ Humberto C Rebelo,⁶¹ Vicente Roqués,⁶² Kristopher M Schroeder,⁶³ Gary Schwartz,⁶⁴ Eric S Schwenk ³, Luc Sermeus ⁶⁵, George Shorten,^{32,33,66} Karthikeyan Srinivasan ¹³, Markus F Stevens,⁶⁷ Kassiani Theodoraki ⁶⁸, Lloyd R Turbitt,⁶⁹ Luis Fernando Valdés-Vilches,⁷⁰ Thomas Volk ^{71,72}, Katrina Webster,^{73,74} T Wiesmann,^{75,76} Sylvia H Wilson ⁷⁷, Morné Wolmarans,⁷⁸ Glenn Woodworth ⁷⁹, Andrew K Worek,⁸⁰ E M Louise Moran ²

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rapm-2021-103136>).

For numbered affiliations see end of article.

Correspondence to

Dr E M Louise Moran, Department of Anaesthesia, Letterkenny University Hospital, Letterkenny F92 AE81, Donegal, Ireland; lmoran75@gmail.com

HMA, BPA and GGC are joint first authors.

Received 27 August 2021
Accepted 10 January 2022



© American Society of Regional Anesthesia & Pain Medicine 2022. Re-use permitted under CC BY-NC. No commercial re-use. Published by BMJ.

To cite: Ahmed HM, Atterton BP, Crowe GG, et al. *Reg Anesth Pain Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/rapm-2021-103136

ABSTRACT

Background and objectives Documentation is important for quality improvement, education, and research. There is currently a lack of recommendations regarding key aspects of documentation in regional anesthesia. The aim of this study was to establish recommendations for documentation in regional anesthesia.

Methods Following the formation of the executive committee and a directed literature review, a long list of potential documentation components was created. A modified Delphi process was then employed to achieve consensus amongst a group of international experts in regional anesthesia. This consisted of 2 rounds of anonymous electronic voting and a final virtual round table discussion with live polling on items not yet excluded or accepted from previous rounds. Progression or exclusion of potential components through the rounds was based on the achievement of strong consensus. Strong consensus was defined as $\geq 75\%$ agreement and weak consensus as 50%–74% agreement.

Results Seventy-seven collaborators participated in both rounds 1 and 2, while 50 collaborators took part in

round 3. In total, experts voted on 83 items and achieved a strong consensus on 51 items, weak consensus on 3 and rejected 29.

Conclusion By means of a modified Delphi process, we have established expert consensus on documentation in regional anesthesia.

INTRODUCTION

Accurate and concise documentation is important for both healthcare delivery and medicolegal protection. Several international medical governing bodies place accurate record keeping as one of the fundamental requirements for good clinical care.^{1,2}

Despite the increase in use of regional anesthesia, there is limited information regarding characteristics of effective documentation. This is in contrast to the wealth of documentation standards which exist for general anesthesia.³

The aim of this study was to establish a consensus opinion among a panel of international experts

regarding the recommended components of documentation in regional anesthesia.

The recommendations contained herein do not define standard of care. They are not intended to replace clinical judgment. In the imperfect setting of heterogeneity of the data, limited data, controversial topics, and bias inherent to expert opinion, compliance with the recommendations may not result in improved outcomes compared with alternative therapies consistent with personalized medicine.

METHODS

An international executive committee consisting of JLB, MJ, HMA, EMLM, and EV was created to design the format of the study. A modified Delphi methodology was chosen as it is a widely used systematic process for achieving consensus amongst a group of experts.⁴ For this study it was decided that a 3 round, prospective Delphi process would be used with 2 rounds of electronic voting and a final round consisting of a round-table discussion and live polling. A steering committee consisting of BPA and GCC was formed to facilitate the management of the Delphi process as well as analyze the results of the voting and communicate with the panel of experts.

Collaborator selection

One-hundred and three experts from North America (n=37), Europe (n=53), and Australasia (n=13) were invited to participate. All those invited were known to have extensive clinical, educational or research experience in the field of regional anesthesia. In addition, a medicolegal expert from the USA was included to advise on any legal implications arising from the consensus document but did not participate in the voting process.

Generation of the long list

A directed literature review was performed to create an exhaustive list of documentation components for performing a regional anesthetic. A MEDLINE search was conducted using the following terms: 'regional anaesthesia documentation', 'peripheral nerve block documentation', 'consent for regional anaesthesia', 'regional anaesthesia litigation', 'documentation', and 'anaesthesia records' yielding 63 potential documentation components. Following review and discussion by the executive committee, an additional 12 items were added to produce the final long list (online supplemental appendix 1).

Modified Delphi process

A modified Delphi methodology was chosen as it is a widely used systematic process for achieving consensus among a group of experts. It is characterized by the generation of a long list of items followed by multiple rounds of anonymous voting and feedback. For this study, all experts who had agreed to take part were invited to participate in 2 electronic voting rounds and a third virtual round table discussion with live polling (figure 1). Strict deadlines were imposed on collaborators to complete each round of the Delphi process within a prescribed timeline. Any collaborators who did not vote within the timeline were excluded from all subsequent rounds and were not included as an author on final publication.

In previous Delphi studies consensus has been defined as $\geq 75\%$ agreement between collaborators.⁵ In this study, 50%–74% was deemed to be weak consensus, while $\geq 75\%$ was agreed to be strong consensus. These ranges were used throughout the Delphi rounds to decide on each item's progression and ultimate inclusion in the final list of suggested documentation components.




Figure 1 Flowchart showing structure of Delphi process and progress of experts throughout the project.




Figure 2 Flowchart showing structure of Delphi process and progress of items throughout the project.

Rounds 1 and 2 were conducted using an online survey tool (Survey Monkey, Momentive, San Mateo, California, USA) distributed via email. Collaborators were asked to ‘agree’ or ‘disagree’ with each of the potential documentation components included in the long list. Commentary and suggestions for additional items were encouraged and were recorded via the same online tool. In round 1, potential documentation components achieving $\geq 50\%$ agreement proceeded to round 2, while those with $< 50\%$ agreement were excluded (figure 2). Following review of the comments made by the collaborators in round 1, some items were reworded or clarified, and any suggested additional items were included for round 2 voting.

In round 2, items achieving $\geq 75\%$ agreement were accepted into the final list of suggested documentation components, while those achieving $< 50\%$ were excluded. Items with $50\% - 74\%$ agreement proceeded to round 3 for further discussion. Based on comments made by collaborators in round 2, some items were reworded or clarified. Any suggested additional items were included for consideration in the round 3 live discussion.

Round 3 was conducted via online videoconferencing software with polling capabilities (Zoom, Zoom Video Communications, San Jose, California, USA). The study included collaborators from multiple countries spanning different time zones, therefore it was unlikely that all collaborators would be available at the same time to take part in the discussion. To decide on a time for round 3 to take place we used online meeting-scheduling software (Doodle, Zurich, Switzerland) to allow collaborators to pick preselected times which would suit their schedule. The most mutually convenient time was picked and circulated to the collaborators via email. Those collaborators unable to attend the live third round were invited to submit comments in advance of the meeting via another survey (Survey Monkey, Momentive, San Mateo, California, USA). A summary of the outcomes of rounds 1 and 2 was also distributed to all collaborators prior to the virtual discussion to avoid confusion about items previously excluded and allow for a more productive discussion. The meeting was co-chaired by EMLM and JLB, who facilitated discussion amongst collaborators via both electronic messaging and oral discourse. Strict adherence to time limits on discussion

(5 minutes) and voting (1 minute) were enforced. All comments made by collaborators in absentia were displayed on-screen for participants during the discussion to ensure all opinions were considered prior to live polling. Items achieving $\geq 75\%$ agreement were accepted into the final list of suggested documentation components. Those achieving $50\% - 74\%$ were accepted as weak agreement, and those achieving $< 50\%$ were excluded.

RESULTS

One-hundred and three experts in regional anaesthesia were invited to take part in the Delphi process. In total 79 agreed to participate (77%); 26 from North America (26/37, 70%), 40 from Europe (40/53, 75%) and 13 from Australasia (13/13, 100%). Seventy-seven collaborators who agreed to take part in the study (97%) completed both rounds 1 and 2. Fifty of the collaborators who originally agreed to take part in the study (65%) were available and took part in the round 3 virtual discussion. The median number of voters for each item in round 3 was 48 (range: 46–50).

Seventy-five potential items were included in round 1 of which 67 achieved $\geq 50\%$ agreement and progressed to round 2. Eight items achieved $< 50\%$ agreement and were excluded (table 1). Responses and collaborator commentary were collected,

Table 1 Items rejected from round 1

Item	% agreement
Patient body mass index	41
Block requested by surgeon	38
Baseline Visual Analog Scale (VAS)	33
Rationale for block performance under spinal, epidural or general anaesthesia	38
Gown used	45
Ultrasound probe decontaminated according to local requirements	42
Needle depth before injection	40
Post-procedure VAS	40

Box 1 Additional items added during rounds 1, 2 and 3**Items added for round 2**

1. Grade of block performer (e.g. Consultant, Fellow, Resident, Registrar)
2. Name of supervisor (if applicable)
3. Name of assistant (if applicable)
4. Method used to secure catheter
5. Ultrasound image included in patient record (if ultrasound used)
6. Post block monitoring completed by ____ / handed over/off to ____ (e.g. Recovery, Post Anesthetic Care Unit (PACU))

Items added for round 3

1. Presence / absence of nerve swelling

Items added during round 3

1. Aseptic technique used as per local policy

anonymized, and summarized by the steering committee and reviewed by the executive committee.

Following review, 15 of the items were reworded or clarified and an additional 6 items were added for round 2 (box 1).

All responses from round 2 were collected and summarized by the steering committee as in round 1. Seventy-three items were included in round 2, of which four did not achieve consensus and were excluded (table 2). Forty-five items achieved $\geq 75\%$ agreement and progressed to the final list of suggested documentation components. The remaining 24 items achieved 50%–74% agreement and were therefore included in round 3 for further discussion. A single additional item was added for round 3 following review of the comments made by the collaborators during round 2 (box 1).

Twenty-five potential items were discussed and voted on in round 3. Five items achieved $\geq 75\%$ agreement and were accepted to the final list of documentation components. Following an ad hoc vote by the collaborators, 4 items pertaining to the use of sterile technique ('sterile gloves used', 'mask used', 'sterile ultrasound transducer cover used' and 'sterile drape used'), were combined into a single all-encompassing item: 'aseptic technique used as per local policy'. This was added to the list of items to be voted on during round 3 and achieved 75% consensus and was duly accepted to the final list of suggested documentation components. Three potential components achieved only 50%–74% agreement and were included in the final list as weak agreement. Thirteen failed to achieve $> 50\%$ agreement and were therefore excluded. It is worth noting that 4 of these were the individual items subsequently grouped together into a single documentation component relating to the use of sterile technique which was accepted.

In total 51 items achieved strong consensus ($\geq 75\%$ agreement), 3 achieved weak consensus (50%–74% agreement) and 29 were rejected ($< 50\%$ agreement). A final list of the suggested

documentation components for regional anesthesia is shown in table 3. A summary of all rounds of voting is provided in online supplemental appendix 2.

DISCUSSION

This study is, to our knowledge, the first to attempt and to achieve an international expert consensus recommendation on documentation in regional anesthesia. Important work has been conducted previously in this area, however only at a national level.^{6,7} It is hoped that our work will facilitate local institutions in their efforts to support physicians with their documentation and quality improvement needs.

While the majority of the potential documentation components were identified in the literature review, several items were added for consideration due to their necessity in billing processes in privately funded healthcare systems. The significant variation in current documentation practices in different countries, in particular between privately and publicly funded healthcare systems, was a theme throughout the study. We encountered commentary in all Delphi rounds that some items required inclusion in certain countries to ensure payment for the performing anesthesiologist. These same items were deemed unnecessary by collaborators from health systems where documentation does not directly impact an individual clinician's reimbursement. This issue is reflected in the voting patterns between the USA versus non-USA anesthesiologists (for full list of USA vs non-USA voting, see online supplemental appendix 3). Taking 'surgeon request for regional anesthesia' as an example, 78% of USA anesthesiologists agreed in round 1 that this item should be included as a documentation component versus just 22% of anesthesiologists located outside of the USA. Ultimately surgeon request for regional anesthesia was rejected in round 1 with only 38% agreement.

We anticipated that collaborators might vote according to individual or local standards of practice rather than specifically the documentation thereof. This was an issue in round 1 and was reflected in the collaborator commentary which contained many comments to the effect of 'I don't use X technique' or 'X technique should no longer be used' or 'X piece of equipment is not available in my hospital'. An example which highlights this issue is the use of nerve stimulators. An item included in round 1 was 'No Evoked Motor Response (EMR) $< ___ \text{mA}$ (when applicable)' which received numerous comments stating that collaborators were not clear what this meant, no longer use nerve stimulators or that the use of nerve stimulators does not improve the safety of nerve blocks. These comments were associated with collaborators voting 'disagree'. Despite these comments and votes this item received sufficient agreement (71%) to progress to round 2. To clarify the item in round 2, we changed the wording to 'No EMR $< ___ \text{mA}$ (when applicable i.e. when nerve stimulator used)'. This item subsequently received 81% agreement in round 2. We clarified several other items with the phrase 'if applicable' or 'if used' and reiterated to the collaborators that the purpose of the study was to define recommended documentation components and not their personal practice.

It was found that a number of collaborators had voted against items being included as they would be found elsewhere in the medical record (e.g. patient date of birth, name, vital signs) and therefore would be an unnecessary duplication. This was mentioned particularly in the context of electronic medical records where a good deal of data entry is automatic rather than by direct human input. These systems, however, are by no means universally used. The executive committee agreed that for the

Table 2 Items rejected from round 2

Item	% agreement
Patient age	32
Name of assistant (if applicable)	42
Method used to secure catheter	45
Ultrasound image included in patient record	42

Table 3 Final list of recommendations

	Level of agreement
Patient information	
Patient name	Strong
Patient date of birth	Strong
Patient gender	Strong
Patient medical record number/hospital number	Strong
Patient weight	Strong
Patient height	Strong
Patient American Society of Anesthesiologists (ASA) physical status classification	Strong
Patient allergies	Strong
Procedure preparation	
Block performed by _____ (name)	Strong
Grade of block performer (e.g. consultant, fellow, resident, registrar)	Weak
Name of supervisor (if applicable)	Strong
Documentation of patient consent gained (as per local standards e.g. written, verbal)	Strong
Documentation of individual risks of procedure discussed (as per local standards)	Strong
Pre-anesthetic / block evaluation	Strong
Coagulation considered	Weak
Pre-procedure diagnosis (post-operative pain management / surgical diagnosis)	Strong
Timeout / World Health Organisation (WHO) checklist	Weak
Stop moment performed	Strong
Intravenous access	Strong
Regional anesthesia procedure name	Strong
Patient position during regional anesthesia procedure	Strong
Monitors applied	Strong
Baseline vital signs	Strong
Pre-medication (type and quantity of sedation)	Strong
Level of sedation (no sedation / light sedation / deep sedation / general anesthesia)	Strong
Procedure performance	
Time and date of regional anesthesia procedure	Strong
Aseptic agent used	Strong
Aseptic technique used as per local policy	Strong
Skin infiltration with local anesthetic	Strong
Needle design: tip, manufacturer, length, gauge	Strong
Local anesthetic used for regional anesthesia technique (concentration and volume)	Strong
Epinephrine dose if used (concentration)	Strong
Adjunct used (e.g. bicarbonate, clonidine etc.)	Strong
Specific for peripheral nerve block performance	
Side of block	Strong
Technique of needle localization (ultrasound / nerve stimulator / landmark)	Strong
No Evoked Motor Response (EMR) < _____ mA (when applicable i.e. when nerve stimulator used)	Strong
Minimum current and current duration (if nerve stimulator used)	Strong
Absence of blood on aspiration	Strong
Catheter depth at the skin	Strong
Absence of pain / paresthesia during injection	Strong
Complications	Strong
Specific for neuraxial procedure performance	
Technique (approach used eg, median/paramedian)	Strong
Vertebral level of needle insertion	Strong
Technique used: loss of resistance to saline/air for epidural insertion	Strong
No of attempts	Strong

Continued

Table 3 Continued

Epidural needle depth at loss of resistance	Strong
Catheter depth at the skin	Strong
Note on aspiration and action taken	Strong
Epidural test dose (if applicable)	Strong
Absence of pain/paresthesia during injection	Strong
Dermatomal level of spinal of epidural block achieved (if assessed)	Strong
Complications	Strong
Postprocedure	
Patient vital signs after the procedure	Strong
Postprocedure instructions (as per local standards)	Strong

purpose of creating an exhaustive list of suggested documentation components the issue of duplication should be ignored and addressed later by individuals or governing bodies designing documentation guidelines. To this end, we asked collaborators to visualize the regional anesthesia record document as a stand-alone 'blank page' in the absence of the rest of the patient's record and other anesthesia documentation. Ultimately a significant number of the suggested items in the finalized list are documented elsewhere in the patient's record independent of a regional anesthesia procedure. Thus, every item may not need to be included on the dedicated regional anesthesia record should it be documented elsewhere.

We have broadly divided our long list and the subsequent finalized list of suggested documentation components into 6 sections.

Patient information

The majority of items relating to patient information were accepted in the early stages of the Delphi process. 'Patient age' was excluded in round 1 on the grounds that 'Patient date of birth' is preferable as, unlike age, it is a unique patient identifier and can be easily used to calculate age if required. 'Patient body mass index' was excluded in round 2 for similar reasons in that it can also be easily calculated.

Procedure preparation

Identification of the block performer by name was accepted to the final list however the grade of the block performer, which was added for round 2, achieved only weak agreement (68%). Many collaborators felt that the name of the block performer would be sufficient information to identify the individual. The name of the supervisor (in the context of a trainee performing a procedure) was added and accepted to the final list in round 2. 'Name of assistant' was added and excluded in round 2 as it was felt that ultimate responsibility lay with the performing or supervising anesthesiologist and therefore the name of any assistants was not essential.

In response to numerous collaborator comments 'Patient consent' and 'Individual risks of procedure discussed/documentation' were reworded in round 2 as 'Documentation of patient consent as per local standards' and 'Documentation of individual risks of procedure discussed (as per local standards)'. It was felt it would be impossible to accurately capture the nuanced aspects of consent internationally, and therefore an open approach was applied to allow scope for local interpretation based on pre-existing standards and legal frameworks. Consent and risks, if discussed, should be documented.

Pre-anesthetic/block evaluation was reworded (from pre-anesthetic evaluation) as collaborators felt that this would be separate to a routine pre-anesthetic assessment and might include

pre-existing neurology or deficits. 'Pre-procedure diagnosis (post-operative pain management/surgical diagnosis)' and 'Indication for regional anesthesia (surgical anesthesia or analgesia)' were considered by many collaborators to be interchangeable and as such only 'Pre-procedure diagnosis (post-operative pain management/surgical diagnosis)' was accepted following discussion in round 3. The recording of the patient's pre-procedure and post-procedure Visual Analog Score (VAS) were rejected in round 1 as collaborators felt that while they might be of research interest, they were of limited clinical relevance.

There was detailed discussion around the inclusion of 'Stop Before You Block (SBYB)' and the World Health Organisation (WHO) Surgical Safety Checklist/time out. The SBYB campaign⁸ originated in the UK and has gained some international acceptance, however, as reported by many collaborators, it is not globally recognized and thus not used in all countries. While the WHO checklist is more widely used, the argument was made that this is performed prior to surgery and not necessarily prior to a block. This is important when regional anesthesia is not performed for surgical anesthesia e.g. labor epidural, or is performed outside the theater e.g. in a dedicated block bay. It was generally agreed that should a 'stop' moment occur it should be documented, however the precise format of this remains unclear and should be determined by local practices.

Consideration of the patient's coagulation status ('Coagulation considered') was accepted with weak agreement. Many collaborators commented that while it should be considered, it was not necessary to document it separately as it forms part of the routine pre-anesthetic/block assessment.

Procedure performance

As previously discussed, 4 items in the original long list relating to aseptic technique were combined into a single item, 'aseptic technique as per local guidelines', which was accepted.

Documentation specific to peripheral nerve block procedures

'Side of Block' was accepted to the final list, however, documentation of 'Block side marked' was excluded in the final roundtable discussion, as it was felt to be repetitive and was adequately addressed elsewhere e.g. by SBYB or a similar stop-moment.

The commentary surrounding the documentation of nerve stimulator technique is perhaps a reflection of changes in regional anesthesia practice; a number of collaborators stated they were not familiar with, or no longer used, several of the techniques described. 'No EMR < ___mA (when applicable i.e. when nerve stimulator used)' and 'Minimum current and current duration (when applicable i.e. when nerve stimulator used)' progressed to the final list. 'Description of quality of paresthesia' and 'Description of motor response' were rejected on the basis that if a nerve locating method was used and a defined end point reached such as paresthesia or muscle contraction, that a description of this was too much detail. 'Catheter tip location confirmed by ultrasound/nerve stimulator' and 'Technique of injection (via needle or catheter)' were also rejected due to perceived excessive detail without providing additional information. 'Note on incremental injection' was rejected on the basis that while incremental injection techniques are advocated, it need not be documented. 'Extra neural spread visualized' and 'Presence/absence of nerve swelling' were also discussed and ultimately rejected, with several collaborators making the point that nerve swelling/extraneural spread themselves were not reliable signs relating to nerve injury. 'Note on resistance to injection < 15 psi' was rejected on the basis that the majority of the expert panel did not have manometry

available nor routinely measure injection pressures with some citing a lack of evidence for its benefit.

Documentation specific to neuraxial nerve block procedures

All but one of the items specific to neuraxial anesthesia were accepted to the final list. 'Method used to secure catheter' was added for round 2 following review of commentary and was rejected. 'Epidural test dose given' and 'Dermatomal level of spinal or epidural block achieved' were clarified in round 2 with 'if applicable', acknowledging that many anesthesiologists choose not to give a test dose, and that dermatomal levels may not be tested prior to the start of surgery or general anesthesia.

Post procedure

'Inclusion of ultrasound image in block note' was added for round 2 based on comments made during round 1. This proved to be somewhat contentious with many remarking that while it would be ideal to have ultrasound images in the patient's record this comes with significant logistical issues, offers no safety benefit and even in medicolegal situations a single static image is of little use. There was a significant divide between USA and non-USA votes on this item, with 82% of USA voters agreeing to this standard in round 2 versus just 25% of non-USA voters. It was ultimately rejected however it will be interesting to see how this evolves in the future; the inclusion of images in the patient's notes has become common practice among laparoscopic surgeons and endoscopists amongst others.

With regard to adequacy of the block for surgery, it was acknowledged that not all blocks are used for surgical anesthesia, and perhaps this item should have been altered to reflect this. Other collaborators made the point that a partial block requiring supplementation is not necessarily a failed block. 'Post-block monitoring completed by ____/ handed over/off to ____ (eg, recovery, PACU)' was rejected as it was felt that this was beyond the scope of the project.

This study has several strengths. Seventy-seven international experts participated in the study exceeding the median number of 17 participants in other Delphi studies reported in the literature.⁴ A large number of experts were deliberately sought to minimize individual bias and provide a broad international perspective on this nuanced area. There was an excellent response rate with all 77 collaborators participating fully in the first 2 rounds and 50 taking part in round 3 despite the scheduling challenges associated with timezones. Voting anonymity was maintained throughout all rounds reducing the impact of dominant individuals, peer pressure, bias and to allow consideration of all opinions in a non-adversarial manner. Collaborator commentary was permitted and encouraged throughout the project allowing for a dynamic long list of suggested documentation components. Items could be added and/or revised based on collaborator commentary; this was well demonstrated during the round 3 live discussion where 4 items were consolidated into a single all-encompassing item in response to real-time collaborator discussion. This flexibility yielded a more comprehensive final long list of suggested documentation components that reflects the varied practice of regional anesthesia world-wide.

This study also has a number of limitations. While the expert panel was international, the majority of the collaborators were from Europe, followed by North America, and with a small number from Australia, New Zealand and Singapore. Our panel did not include any collaborators from South America, the Middle East, Asia or Africa, nor were there representatives from low- and middle-income countries. Thus, our suggested

documentation components may not be applicable in these contexts. A limitation of all Delphi studies is that any expert panel will be influenced by their own experience and personal practice, however as previously mentioned, the large number of experts in this project should minimize this effect. Another potential limitation arises from the small number of prior studies in this area. As a result, the long list formulation required significant input from the executive committee to generate what was felt to be complete list of documentation components. Twelve additional items were added to the long list originally derived from the literature review, and a further 8 were added during the Delphi process, which could represent a source of bias.

The practice of regional anesthesia is constantly evolving and as such this list of suggested documentation components requires regular review to remain current. It is hoped that this list might provide a framework for international regional anesthesia societies to produce guidelines for documentation standards in the near future. There is no single global standard of care. The practice of medicine and regional anesthesia vary dramatically across the world and documentation should rightly reflect local best practice.

Concerning areas for future research, these may include the formation of a standard block procedure note template for both electronic and paper medical records and standardizing post-block monitoring and follow-up documentation for regional anesthesia procedures including inpatient and ambulatory nerve catheters. Another evolving area that warrants future study is the utility of including digital ultrasound media within the patient's record should this practice become more widespread.

The authors acknowledge that while attempts to improve documentation standards are necessary and well-intended, an excessive burden of notes may distract from patient care and can even lead to physician burnout.⁹ Although the final list of recommendations may seem extensive the majority will already be included in the routine documentation practice of most anesthesiologists. This is not intended to be an exhaustive list nor a legal standard of documentation but rather a consensus of useful items to document patient care.

CONCLUSION

By means of a modified Delphi process we have established an expert panel consensus on documentation in regional anesthesia. We hope adoption of our recommendations will facilitate physician workflow, education, quality improvement and research.

Author affiliations

- ¹Department of Anaesthesia, Leeds Teaching Hospitals NHS Trust, Leeds, UK
- ²Department of Anaesthesia and Intensive Care, Letterkenny University Hospital, Letterkenny, Donegal, Ireland
- ³Department of Anesthesiology, Thomas Jefferson University Sidney Kimmel Medical College, Philadelphia, Pennsylvania, USA
- ⁴Department of Anaesthesia, Fiona Stanley Hospital and Freemantle Hospitals, Perth, Western Australia, Australia
- ⁵Department of Anesthesiology and Perioperative Medicine, Penn State College of Medicine, Hershey, Pennsylvania, USA
- ⁶Department of Anaesthesia, University Hospital of Lausanne, Lausanne, Switzerland
- ⁷Faculty of Biology and Medicine, University of Lausanne, Lausanne, Switzerland
- ⁸Department of Anesthesiology, Critical Care and Pain Medicine, Boston Children's Hospital, Boston, Massachusetts, USA
- ⁹Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University, Palo Alto, California, USA
- ¹⁰Department of Anaesthesia, Intensive Care and Pain Medicine, St Vincent's University Hospital, Dublin, Ireland
- ¹¹Department of Anaesthesia, Waikato Hospital, Hamilton, New Zealand
- ¹²Department of Anaesthesia, Liverpool Hospital, University of New South Wales Faculty of Medicine, Sydney, New South Wales, Australia
- ¹³Department of Anaesthesia, Tallaght University Hospital, Dublin, Ireland
- ¹⁴Department of Anaesthesia and Pain Medicine, Gold Coast University Hospital, Gold Coast, Queensland, Australia

- ¹⁵Faculty of Medical Sciences, Unit of Pain Research and Treatment, Medical University of Silesia, Zabrze, Poland
- ¹⁶Department of Anaesthesia, Clinique Medipole Garonne, Toulouse, France
- ¹⁷Department of Anesthesia and Intensive Care, Cluj-Napoca County Emergency Hospital, Cluj-Napoca, Romania
- ¹⁸Department of Anaesthesiology, Intensive Care and Pain Medicine, National Pirogov Memorial Medical University, Vinnytsia, Ukraine
- ¹⁹Department of Anaesthesia and Pain Management, Royal Children's Hospital Melbourne, Melbourne, Victoria, Australia
- ²⁰Retired Consultant Anaesthetist, Worcester Royal Hospital, Worcester, UK
- ²¹Department of Anaesthesia, Mater Misericordiae University Hospital, Dublin, Ireland
- ²²Department of Anesthesiology, Duke University Medical Center, Durham, North Carolina, USA
- ²³Department of Anesthesia, Clinique Sainte Anne Saint Remi, Bruxelles, Belgium
- ²⁴Department of Anesthesiology, Vanderbilt University Medical Center, Nashville, Tennessee, USA
- ²⁵Department of Anesthesiology and Reanimation, Koç University Hospital, Istanbul, Turkey
- ²⁶Department of Anesthesiology, University of North Carolina School of Medicine, Chapel Hill, North Carolina, USA
- ²⁷Department of Anaesthesia, St Mary's Hospital, Imperial College, London, UK
- ²⁸Ultrasound Education Unit, University of Melbourne, Melbourne, Victoria, Australia
- ²⁹Department of Anaesthesia, Northeast Health Wangaratta, Wangaratta, Victoria, Australia
- ³⁰Department of Anesthesiology, University of Texas McGovern Medical School, Houston, Texas, USA
- ³¹Department of Anaesthesiology and Intensive Care, F D Roosevelt Teaching Hospital, Banska Bystrica, Slovakia
- ³²Department of Anaesthesiology and Intensive Care Medicine, Cork University Hospital, Cork, Ireland
- ³³School of Medicine, University College Cork, Cork, Ireland
- ³⁴Department of Anesthesia and Pain Medicine, University of Alberta Hospital, Edmonton, Alberta, Canada
- ³⁵Department of Anesthesiology, Perioperative and Pain Medicine and Orthopaedics, Icahn School of Medicine at Mount Sinai, New York, New York, USA
- ³⁶Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, Minnesota, USA
- ³⁷Department of Anesthesiology and Perioperative Medicine, The University of Alabama, Birmingham, Alabama, USA
- ³⁸Department of Anaesthesia, Galway University Hospitals, Galway, Ireland
- ³⁹Department of Anaesthesia, Royal Prince Alfred Hospital, Sydney, New South Wales, Australia
- ⁴⁰Anesthesiology and Perioperative Care Service, Veterans Affairs Palo Alto Health Care System, Palo Alto, California, USA
- ⁴¹Department of Anesthesiology, Perioperative and Pain Medicine, Stanford Health Care, Stanford, California, USA
- ⁴²Department of Anesthesia and Surgical Intensive Care, Changi General Hospital, Singapore
- ⁴³Department of Anaesthesia, Hospital das Forças Armadas, Porto, Portugal
- ⁴⁴Department of Anesthesiology, Columbia University Irving Medical Center, New York, New York, USA
- ⁴⁵Department of Anaesthesia, Pain and Critical Care Medicine, Glasgow Royal Infirmary, Glasgow, UK
- ⁴⁶School of Medicine, University of Glasgow, Glasgow, UK
- ⁴⁷Department of Anesthesiology and Pain Management, University of Texas Southwestern Medical Center, Dallas, Texas, USA
- ⁴⁸Department of Anaesthesia, South Infirmary Victoria University Hospital, Cork, Ireland
- ⁴⁹Department of Anaesthesia, Flinders Medical Centre, Adelaide, South Australia, Australia
- ⁵⁰Department of Anaesthesia, Craigavon Area University Teaching Hospital, Craigavon, UK
- ⁵¹Department of Anaesthesiology, Hospital of Traumatology and Orthopaedics, University of Latvia, Riga, Latvia
- ⁵²Department of Anaesthesia, Sir Charles Gairdner Hospital, Perth, Western Australia, Australia
- ⁵³Department of Anaesthesiology, Creta InterClinic Hospital, Hellenic Healthcare Group, Heraklion-Crete, Greece
- ⁵⁴Department of Anesthesiology, University of Florida College of Medicine, Gainesville, Florida, USA
- ⁵⁵Department of Theatres, Anaesthesia and Perioperative Medicine, Guy's and St Thomas' NHS Foundation Trust, London, UK
- ⁵⁶Faculty of Life Sciences and Medicine, King's College London, London, UK
- ⁵⁷Department of Anesthesia and Pain Medicine, Toronto Western Hospital, Toronto, Ontario, Canada
- ⁵⁸Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada
- ⁵⁹Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Jacksonville, Florida, USA

- ⁶⁰Department of Anesthesiology, University of Kansas Health System, Kansas City, Kansas, USA
- ⁶¹Department of Anaesthesiology, Hospital Luz Arrábida, Vila Nova de Gaia, Portugal
- ⁶²Department of Anaesthesia and Intensive Care Medicine, Virgen de la Arrixaca University Hospital, Murcia, Spain
- ⁶³Department of Anesthesiology, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, USA
- ⁶⁴Department of Anesthesiology, Maimonides Medical Center, Brooklyn, New York, USA
- ⁶⁵Department of Anesthesia, Cliniques Universitaires Saint-Luc, UCLouvain, Bruxelles, Belgium
- ⁶⁶Insight II, SFI Research Centre, Cork, Ireland
- ⁶⁷Department of Anesthesiology, Amsterdam University Medical Centers, Amsterdam, Netherlands
- ⁶⁸Department of Anesthesiology, National and Kapodistrian University of Athens, Aretaieion University Hospital, Athens, Greece
- ⁶⁹Department of Anaesthesia, Royal Victoria Hospital, Belfast, UK
- ⁷⁰Department of Anesthesiology and Reanimation, Hospital Costa del Sol, Marbella, Spain
- ⁷¹Department of Anaesthesiology, Intensive Care and Pain Therapy, Saarland University Hospital, Homburg, Germany
- ⁷²Faculty of Medicine, Saarland University, Homburg, Germany
- ⁷³Department of Anaesthesia and Perioperative Medicine, Royal Hobart Hospital, Hobart, Tasmania, Australia
- ⁷⁴School of Medicine, University of Tasmania, Hobart, Tasmania, Australia
- ⁷⁵Department of Anesthesiology & Intensive Care Medicine, University Hospitals Giessen and Marburg Campus Giessen, Giessen, Germany
- ⁷⁶Department of Anesthesiology and Intensive Care Medicine, Diakoneo Diak Klinikum, Schwäbisch Hall, Germany
- ⁷⁷Department of Anesthesia and Perioperative Medicine, Medical University of South Carolina, Charleston, South Carolina, USA
- ⁷⁸Department of Anaesthesia, Norfolk and Norwich University Hospital NHS Trust, Norwich, UK
- ⁷⁹Department of Anesthesiology and Perioperative Medicine, Oregon Health and Science University, Portland, Oregon, USA
- ⁸⁰Post and Post LLC, Berwyn, Pennsylvania, USA

Twitter Hassan M Ahmed @HassanM31118046, Sanjib Adhikary @sanjibadhikary, Eric Albrecht @DrEAlbrecht, Jan Boublik @janboublik MDPHd, Alwin Chuan @AlwinChuan, Dan Sebastian Dirzu @DanDirzu, Rajnish K Gupta @dr_rajgupta, Harold David Hardman @hdhardman_david, Jakub Hlasny @HlasnyJakub, Rebecca L Johnson @rljohnsonmd, Andrew Kenneth Lansdown @TheLansdowns, Clara Lobo @claralexlobo, Alan James Robert Macfarlane @ajrmacfarlane, Peter Merjavý @PeterMerjavý, Peter Moran @PetermoranPeter, Amit Pawa @amit_pawa, Anahi Perlas @PerlasAnahi, Kristopher M Schroeder @KristopherSch6, Gary Schwartz @garyschwartzmd, Eric S Schwenk @ESchwenkMD and E M Louise Moran @LouiseMoran75

Acknowledgements This paper is supported by the ASRA Pain Medicine and the ESRA board of directors.

Contributors Study conception and design: HMA, BPA, GGC, JLB, MJ, EV, EMLM and AKW. Study conduct: HMA, BPA, GGC, JLB, MJ, EV and EMLM. Data analysis: BPA and GGC. Manuscript preparation: HMA, BPA, GGC, JLB, MJ, EV and EMLM. Manuscript revision: all authors. Manuscript approval: all authors. Manuscript guarantor: EMLM

Funding The authors received a grant towards the cost of open access from The European Society of Regional Anaesthesia and Pain Therapy

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Institutional ethics committee approval was deemed unnecessary by Letterkenny University Hospital Ethics Committee in December, 2020.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Any data not in the online supplemental information are available on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, an indication of whether changes were made, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Hassan M Ahmed <http://orcid.org/0000-0001-5328-6039>
 Mark Johnson <http://orcid.org/0000-0002-1503-8955>
 Eugene Viscusi <http://orcid.org/0000-0003-0260-4396>
 Sanjib Adhikary <http://orcid.org/0000-0001-7167-1184>
 Eric Albrecht <http://orcid.org/0000-0001-6432-1311>
 Kelly Byrne <http://orcid.org/0000-0002-7890-7265>
 Alwin Chuan <http://orcid.org/0000-0003-4356-6525>
 Dan Sebastian Dirzu <http://orcid.org/0000-0002-6730-7045>
 Dmytro Dmytriiev <http://orcid.org/0000-0001-6067-681X>
 Rajnish K Gupta <http://orcid.org/0000-0003-3401-4737>
 Vivian H Y Ip <http://orcid.org/0000-0001-6158-4415>
 Christina L Jeng <http://orcid.org/0000-0003-2507-7015>
 Rebecca L Johnson <http://orcid.org/0000-0002-1920-9774>
 Yean Chin Lim <http://orcid.org/0000-0003-3679-7173>
 Alan James Robert Macfarlane <http://orcid.org/0000-0003-3858-6468>
 Anthony T Machi <http://orcid.org/0000-0001-8245-795X>
 Peter Merjavý <http://orcid.org/0000-0001-6813-3641>
 Amit Pawa <http://orcid.org/0000-0002-2404-9162>
 Anahi Perlas <http://orcid.org/0000-0002-8190-8314>
 Steven Porter <http://orcid.org/0000-0002-8657-2004>
 Eric S Schwenk <http://orcid.org/0000-0003-3464-4149>
 Luc Sermeus <http://orcid.org/0000-0003-3191-430X>
 Karthikeyan Srinivasan <http://orcid.org/0000-0003-2833-1920>
 Kassiani Theodoraki <http://orcid.org/0000-0003-4890-4642>
 Thomas Volk <http://orcid.org/0000-0001-5175-7159>
 Sylvia H Wilson <http://orcid.org/0000-0003-4747-6009>
 Glenn Woodworth <http://orcid.org/0000-0002-1924-801X>
 E M Louise Moran <http://orcid.org/0000-0001-8069-5786>

REFERENCES

- 1 Good Medical Practice. General medical Council. Available: https://www.gmc-uk.org/-/media/documents/good-medical-practice-english-20200128_pdf-51527435.pdf
- 2 Good medical practice: a code of conduct for doctors in Australia. Medical Board of Australia. Available: <https://www.medicalboard.gov.au/codes-guidelines-policies/code-of-conduct.aspx> [Accessed Mar 2021].
- 3 Klein AA, Meek T, Allcock E, *et al*. Recommendations for standards of monitoring during anaesthesia and recovery 2021: guideline from the association of anaesthetists. *Anaesthesia* 2021;76:1212–23.
- 4 Boulkedid R, Abdoul H, Loustau M, *et al*. Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One* 2011;6:e20476.
- 5 Diamond IR, Grant RC, Feldman BM, *et al*. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401–9.
- 6 Moran PJ, Fennessy P, Johnson MZ. Establishing a new national standard for the documentation of regional anaesthesia in Ireland. *BMJ Open Qual* 2017;6:e000210.
- 7 Gerancher JC, Viscusi ER, Liguori GA, *et al*. Development of a standardized peripheral nerve block procedure note form. *Reg Anesth Pain Med* 2005;30:67–71.
- 8 Safe Anaesthesia Liaison Group, Regional Anaesthesia UK. *Stop before you block campaign: supporting information*, 2015.
- 9 Downing NL, Bates DW, Longhurst CA. Physician burnout in the electronic health record era: are we ignoring the real cause? *Ann Intern Med* 2018;169:50–1.

APPENDIX 1.0 - LONG LIST OF SUGGESTED ITEMS FOR DOCUMENTATION IN REGIONAL ANESTHESIA**Patient information:**

1. Name¹⁻⁶
2. Age³⁻⁶
3. Date of birth
4. Gender³
5. Medical record/ hospital number¹⁻⁶
6. Weight^{1,3}
7. Height³
8. BMI
9. ASA^{1,3}
10. Allergy^{1,3}

Procedure Preparation:

11. Block performed by _____ (name)^{1,3,5-8}
12. Consent^{5,7,9}
13. Individual risks of procedure discussed and documented^{1,9,10}
14. Pre-anesthetic evaluation⁷
15. Pre-op diagnosis (Post-operative pain management / surgical diagnosis)^{5,6}
16. Timeout / WHO checklist^{5,6}
17. Stop moment performed
18. Coagulation considered⁵
19. IV access⁵
20. Block procedure name^{1,3,5,6}
21. Indication for regional anesthesia (purpose of the block)^{6,8}
22. Request by surgeon for placement for pain management^{6,8}
23. Patient position during the regional anesthesia procedure^{3,6,7}
24. Monitors applied^{2,5,6,11}
25. Baseline vital signs¹²
26. Baseline visual analogue score (VAS)¹³
27. Pre-medication (type and quantity of sedation)^{1,5,6,14,15}
28. Level of sedation (no sedation / light sedation / deep sedation / GA)⁶
29. Indication for block under spinal, epidural or general anesthesia⁵

Procedure Performance:

1. Time and date of regional anesthetic procedure^{6,8,14}
2. Aseptic agent used^{5,6,16}
3. Sterile gloves used
4. Mask used
5. Gown used
6. Ultrasound probe decontaminated in accordance with local policy
7. Sterile ultrasound transducer cover used
8. Sterile drape used⁶
9. Skin infiltration with local anesthetic⁶
10. Needle design: tip, manufacturer, length, gauge^{5,6,17}

11. Local anesthetic used for regional anesthetic technique (concentration and volume)⁵⁻⁷
12. Adrenaline dose used (concentration)⁶
13. Adjunct used^{5,6,18}

Specific for Regional Nerve Block Procedures:

30. Side of block¹⁹
31. Block side marked
32. Technique of needle localization (ultrasound / nerve stimulator / landmark technique)^{5-7,20}
33. Approach used (In plane / out of plane)^{1,3,5,6,9,20}
34. No EMR < ___mA when applicable^{5,6}
35. Minimum current and current duration^{6,21}
36. Description of motor response²⁰
37. Description of quality of paresthesia²²
38. Catheter tip location confirmed by ultrasound / nerve stimulator⁶
39. Technique of injection (via needle or catheter)^{6,23}
40. Note on incremental injection¹¹
41. Extra-neural spread visualized^{5,6}
42. Note on resistance during injection (pressure < 15psi)^{5,6,24}
43. Absence of blood on aspiration^{5,6}
44. Needle depth before injection^{6,21}
45. Catheter depth at the skin^{6,23}
46. Absence of pain / paresthesia during injection^{5,6,14,21}

Specific for Neuraxial Procedures:

47. Approach used (e.g. median / paramedian)^{1,3,5,6,9,20}
48. Vertebral level of needle insertion¹⁴
49. Technique used: Loss of resistance to saline / air for epidural insertion
50. Number of attempts⁵
51. Epidural needle depth at loss of resistance^{6,21}
52. Catheter depth at the skin^{6,23}
53. Note on aspiration and action taken¹¹
54. Absence of pain / paraesthesia during injection^{5,6,14,21}
55. Epidural test dose
56. Dermatomal level of spinal or epidural block achieved^{7,14}
57. Complications⁷

Post Procedure:

58. Adequacy of regional anesthesia procedure^{5,15}
59. Complications⁷
60. Patient vital signs after the procedure²⁵
61. Patient visual analogue score after the procedure¹⁵
62. Post procedure instructions⁵

REFERENCES:

1. Elhalawani I, Jenkins S, Newman N. Perioperative anesthetic documentation: Adherence to current Australian guidelines. *J Anaesthesiol Clin Pharmacol*. 2013;29(2):211-215.
2. Kendall F. Documenting local anesthesia patient care. Developing the tool. *AORN J*. 1993;58(4):715-719.
3. Jang J, Yu SH, Kim CB, Moon Y, Kim S. The effects of an electronic medical record on the completeness of documentation in the anesthesia record. *Int J Med Inform*. 2013;82(8):702-707.
4. The Joint Commission announces the 2005 National Patient Safety Goals and Requirements. Official publication of revised national patient safety goals. *Jt Comm Perspect* 2004;24:8.
5. Moran PJ, Fennessy P, Johnson MZ. Establishing a new national standard for the documentation of regional anaesthesia in Ireland. *BMJ Open Qual*. 2017;6(2):e000210.
6. Gandhi K, Patel V, Maliakal T, Xu D, Flisinski K. Universal Documentation Sheet for Peripheral Nerve Blocks. *The Journal of New York School of Regional Anaesthesia*. 2009;30(1):23-24.
7. Wilbanks BA, Geisz-Everson M, Boust RR. The role of documentation quality in anesthesia-related closed claims: A descriptive qualitative study. *CIN - Comput Informatics Nurs*. 2016;34(9):406-412.
8. Rosenquist RW, Williams BA. Optimizing Billing and Compliance. *Syllabus from the 2004 Annual Spring Meeting of the American Society of Regional Anesthesia and Pain Medicine*, Orlando FL. Pages 333–341
9. Australian And New Zealand College of Anaesthetists. Review Ps26 (2005) Guidelines on Consent for Anaesthesia or Sedation. Vol 26.; 2005.
10. Studdert DM, Mello MM, Levy MK, et al. Geographic Variation in Informed Consent Law : Two Standards for Disclosure of Treatment Risks. *Journal of Empirical Legal Studies*. 2007;4(1):103-124.
11. Mulroy MF. Systemic toxicity and cardiotoxicity from local anesthetics: incidence and preventive measures. *Reg Anesth Pain Med*. 2002;27(6):556-561.
12. Gerancher JC, Viscusi ER, Liguori GA, et al. Development of a standardized peripheral nerve block procedure note form. *Reg Anesth Pain Med*. 2005;30(1):67-71.
13. ASA Standards for Basic Anesthetic Monitoring, Guidelines for Regional Anesthesia in Obstetrics, and Documentation of Anesthesia Care. ASA House of Delegates 1986-2003. Updated 2010. Reaffirmed on December 13, 2020. <https://www.asahq.org/standards-and-guidelines/standards-for-basic-anesthetic-monitoring>
14. Miu M, Paech MJ. Documentation of regional anaesthesia for caesarean delivery. *Anaesth Intensive Care*. 2013;41(2):271-272.
15. Wu CL, Naqibuddin M, Fleisher LA. Measurement of patient satisfaction as an outcome of regional anesthesia and analgesia: a systematic review. *Reg Anesth Pain Med*. 2001;26(3):196-208.
16. Birnbach DJ, Meadows W, Stein DJ, Murray O, Thys DM, Sordillo EM. Comparison of povidone iodine and DuraPrep, an iodophor-in-isopropyl alcohol solution, for skin disinfection prior to epidural catheter insertion in parturients. *Anesthesiology*. 2003;98(1):164-169.
17. Cousins MJ, Bridenbaugh PO, Carr DB, Horlocker TT., *Neural Blockade in Clinical Anesthesia and Management of Pain*. 4th ed. Lippincott Williams & Wilkins (LWW). 2008.

18. Murphy DB, McCartney CJ, Chan VW. Novel analgesic adjuncts for brachial plexus block: a systematic review. *Anesth Analg*. 2000;90(5):1122-1128.
19. Wrong Site Surgery Summit addresses current problems, future solutions. *Jt Comm Perspect* 2003;23:8-9.
20. Neal JM, Hebl JR, Gerancher JC, Hogan QH. Brachial plexus anesthesia: essentials of our current understanding. *Reg Anesth Pain Med*. 2002;27(4):402-428.
21. Hadzic A, Hadzic's *Peripheral Nerve Blocks and Anatomy for Ultrasound-Guided Regional Anesthesia*. McGraw-Hill Medical. 2nd edition; 2012.
22. Lee LA, Posner KL, Domino KB, Caplan RA, Cheney FW. Injuries associated with regional anesthesia in the 1980s and 1990s: a closed claims analysis. *Anesthesiology*. 2004;101(1):143-152.
23. Liu SS, Salinas F V. Continuous plexus and peripheral nerve blocks for postoperative analgesia. *Anesth Analg*. 2003;96(1):263-272.
24. Hadzic A, Dilberovic F, Shah S, et al. Combination of intraneural injection and high injection pressure leads to fascicular injury and neurologic deficits in dogs. *Reg Anesth Pain Med*. 2004;29(5):417-423.
25. McGrath B, Chung F. Postoperative recovery and discharge. *Anesthesiol Clin North America*. 2003;21(2):367-386.

APPENDIX 2.0 – SUMMARY OF ROUNDS 1, 2 AND 3

Item	Round 1	Round 2	Round 3	Overall Outcome
Patient name	Progressed to round 2	Accepted		Accepted
Patient age	Progressed to round 2 and item clarified with “(if you believe date of birth is sufficient, please disagree)”	Rejected		Rejected
Patient date of birth	Progressed to round 2	Accepted		Accepted
Patient gender	Progressed to round 2	For discussion in round 3	78% agree (36/46)	Accepted
Patient medical record number / hospital number	Progressed to round 2	Accepted		Accepted
Patient weight	Progressed to round 2	Accepted		Accepted
Patient height	Progressed to round 2	For discussion in round 3	96% agree (45/47)	Accepted
Patient BMI	Rejected			Rejected
Patient ASA physical status classification	Progressed to round 2	Accepted		Accepted
Patient allergies	Progressed to round 2	Accepted		Accepted
Block performed by _____ (name)	Progressed to round 2	Accepted		Accepted
Grade of block performer (e.g. Consultant, Fellow, Resident, Registrar etc.)	Added for round 2	For discussion in round 3	68% agree (32/47)	Accepted with weak agreement
Name of supervisor (if applicable)	Added for round 2	Accepted		Accepted
Name of assistant (if applicable)	Added for round 2	Rejected		Rejected
Patient consent	Progressed to round 2 and re-worded as “Documentation of patient consent gained (as per local standards e.g. written, verbal)”	Accepted		Accepted
Individual risks of procedure discussed and documented	Progressed to round 2 and re-worded as: “Documentation of individual risks of procedure discussed (as per local standards)”	Accepted		Accepted
Pre-anesthetic evaluation	Progressed to round 2 and re-worded as “Pre-anaesthetic/block evaluation”	For discussion in round 3	96% agree (45/47)	Accepted
Pre-procedure diagnosis (post-operative pain management / surgical diagnosis)	Progressed to round 2	For discussion in round 3	77% agree (36/47)	Accepted
Timeout / WHO checklist	Progressed to round 2	For discussion in round 3	57% agree (28/49)	Accepted with weak agreement
Stop moment performed	Progressed to round 2	For discussion in round 3	80% agree (40/50)	Accepted
Coagulation considered	Progressed to round 2	For discussion in round 3	50% agree (24/48)	Accepted with weak agreement
Intravenous access	Progressed to round 2	Accepted		Accepted

Regional anesthesia procedure name	Progressed to round 2	Accepted		Accepted
Indication for regional anesthesia (surgical anesthesia or analgesia)	Progressed to round 2	For discussion in round 3	48% agree (23/48)	Rejected
Request by surgeon	Rejected			Rejected
Patient position during regional anesthesia procedure	Progressed to round 2	Accepted		Accepted
Monitors applied	Progressed to round 2	Accepted		Accepted
Baseline vital signs	Progressed to round 2	Accepted		Accepted
Baseline visual analogue score	Rejected			Rejected
Pre-medication (type and quantity of sedation)	Progressed to round 2	Accepted		Accepted
Level of sedation (no sedation / light sedation / deep sedation / general anesthesia)	Progressed to round 2	Accepted		Accepted
Rationale for block performance under spinal, epidural or general anesthesia	Rejected			Rejected
Time and date of regional anesthesia procedure	Progressed to round 2	Accepted		Accepted
Aseptic agent used	Progressed to round 2	Accepted		Accepted
Sterile gloves used	Progressed to round 2 and item clarified with "(if applicable)"	For discussion in round 3	6% agree (3/47)	Rejected
Mask used	Progressed to round 2 and item clarified with "(if applicable)"	For discussion in round 3	6% agree (3/47)	Rejected
Gown used	Rejected			Rejected
Ultrasound probe decontaminated according to local requirements	Rejected			Rejected
Sterile ultrasound transducer cover used	Progressed to round 2 and item clarified with "(if applicable)"	For discussion in round 3	2% agree (1/48)	Rejected
Sterile drape used	Progressed to round 2 and item clarified with "(if applicable)"	For discussion in round 3	0% agree (0/47)	Rejected
Aseptic technique used as per local policy			Added during round 3 during live discussion 75% agree (36/48)	Accepted
Skin infiltration with local anesthetic	Progressed to round 2	Accepted		Accepted
Needle design: tip, manufacturer, length, gauge	Progressed to round 2	Accepted		Accepted

Local anesthetic used for regional anesthesia technique (concentration and volume)	Progressed to round 2	Accepted		Accepted
Adrenaline dose if used (concentration)	Progressed to round 2	Accepted		Accepted
Adjunct used	Progressed to round 2 and item clarified with "(e.g. bicarbonate, clonidine etc.)"	Accepted		Accepted
Side of block	Progressed to round 2	Accepted		Accepted
Blocked side marked	Progressed to round 2	For discussion in round 3	28% agree (13/47)	Rejected
Technique of needle localization (ultrasound / nerve stimulator / landmark)	Progressed to round 2	Accepted		Accepted
Approach used (in plane / out of plane)	Progressed to round 2	For discussion in round 3	19% agree (9/47)	Rejected
No EMR < ___mA (when applicable)	Progressed to round 2 and re-worded as "No Evoked Motor Response < ___mA (when applicable i.e. when nerve stimulator used)"	Accepted		Accepted
Minimum current and current duration	Progressed to round 2 and item clarified with "(if nerve stimulator used)"	Accepted		Accepted
Description of motor response	Progressed to round 2 and item clarified with "(if nerve stimulator used)"	For discussion in round 3	17% agree (8/47)	Rejected
Description of quality of paresthesia	Progressed to round 2 and item clarified with "(if using paresthesia technique)"	For discussion in round 3	6% agree (3/48)	Rejected
Catheter tip location confirmed by ultrasound / nerve stimulator	Progressed to round 2	For discussion in round 3	4% agree (2/48)	Rejected
Technique of injection (via needle or catheter)	Progressed to round 2	For discussion in round 3	17% agree (8/48)	Rejected
Note on incremental injection	Progressed to round 2	For discussion in round 3	8% agree (4/48)	Rejected
Extra-neural spread visualized	Progressed to round 2	For discussion in round 3	21% agree (10/48)	Rejected
Presence / absence nerve swelling		Added for round 3	19% agree (9/48)	Rejected
Note on resistance during injection (pressure < 15psi)	Progressed to round 2	For discussion in round 3	8% agree (4/48)	Rejected
Absence of blood on aspiration	Progressed to round 2	Accepted		Accepted
Needle depth before injection	Rejected			Rejected
Catheter depth at the skin	Progressed to round 2	Accepted		Accepted
Absence of pain / paresthesia during injection	Progressed to round 2	Accepted		Accepted
Technique (approach used e.g. median / paramedian)	Progressed to round 2	Accepted		Accepted
Vertebral level of needle insertion	Progressed to round 2	Accepted		Accepted

Technique used: Loss of resistance to saline/air for epidural insertion	Progressed to round 2	Accepted		Accepted
Number of attempts	Progressed to round 2	Accepted		Accepted
Epidural needle depth at loss of resistance	Progressed to round 2	Accepted		Accepted
Catheter depth at the skin	Progressed to round 2	Accepted		Accepted
Method used to secure catheter	Added for round 2	Rejected		Rejected
Note on aspiration and action taken	Progressed to round 2	Accepted		Accepted
Epidural test dose	Progressed to round 2 and item clarified with "(if applicable)"	Accepted		Accepted
Absence of pain / paresthesia during injection	Progressed to round 2	Accepted		Accepted
Dermatomal level of spinal or epidural block achieved	Progressed to round 2 and item clarified with "(if assessed)"	Accepted		Accepted
Complications (neuraxial)	Progressed to round 2	Accepted		Accepted
Adequacy of regional anesthesia procedure	Progressed to round 2	For discussion in round 3	36% agree (17/47)	Rejected
Complications (nerve block)	Progressed to round 2	Accepted		Accepted
Ultrasound image included in patient record (if US used)	Added for round 2	Rejected		Rejected
Patient vital signs after the procedure	Progressed to round 2	Accepted		Accepted
Post block monitoring completed by _____/ handed over/off to _____ (e.g. Recovery, PACU)	Added for round 2	For discussion in round 3	6% agree (3/47)	Rejected
Patient visual analogue score after the procedure	Rejected			Rejected
Post procedure instructions	Progressed to round 2 and item clarified with "(as per local standards)"	Accepted		Accepted

APPENDIX 3.0 – SUMMARY OF US vs NON-US VOTING – ROUNDS 1, 2 AND 3

Item	Round 1 (% agree)			Round 2 (% agree)			Round 3 (% agree)			Overall Outcome
	US	Non-US	Total	US	Non-US	Total	US	Non-US	Total	
Patient name	100%	100%	100%	100%	95%	96%				Accepted
Patient age	78%	53%	60%	41%	29%	32%				Rejected
Patient date of birth	87%	78%	81%	91%	93%	92%				Accepted
Patient gender	78%	73%	74%	77%	69%	71%	84%	74%	78%	Accepted
Patient medical record / hospital number	100%	93%	95%	95%	95%	95%				Accepted
Patient weight	87%	78%	81%	95%	89%	91%				Accepted
Patient height	61%	56%	58%	59%	64%	62%	95%	96%	96%	Accepted
Patient BMI	39%	42%	41%							Rejected
Patient ASA physical status classification	83%	80%	82%	68%	82%	78%				Accepted
Patient allergies	91%	91%	91%	86%	98%	95%				Accepted
Block performed by _____ (name)	100%	96%	97%	100%	100%	100%				Accepted
Grade of block performer (e.g. Consultant, Fellow, Resident, Registrar etc.)				59%	69%	66%	68%	68%	68%	Accepted with weak agreement
Name of supervisor (if applicable)				100%	84%	88%				Accepted
Name of assistant (if applicable)				55%	36%	42%				Rejected
Documentation of patient consent gained (as per local standards e.g. written, verbal)	91%	91%	91%	95%	95%	95%				Accepted
Documentation of individual risks of procedure discussed (as per local standards)	87%	85%	86%	77%	80%	79%				Accepted
Pre-anesthetic / block evaluation	74%	73%	73%	82%	71%	74%	100%	93%	96%	Accepted
Pre-procedure diagnosis (post-operative pain)	87%	71%	76%	91%	64%	71%	84%	71%	77%	Accepted

management / surgical diagnosis)										
Timeout / WHO checklist	96%	67%	76%	95%	65%	74%	90%	35%	57%	Accepted with weak agreement
Stop moment performed	39%	85%	72%	41%	84%	71%	60%	93%	80%	Accepted
Coagulation considered	83%	75%	77%	73%	73%	73%	42%	55%	50%	Accepted with weak agreement
Intravenous access	70%	82%	78%	73%	84%	81%				Accepted
Regional anesthesia procedure name	100%	100%	100%	100%	100%	100%				Accepted
Indication for regional anesthesia (surgical anesthesia or analgesia)	91%	58%	68%	86%	55%	64%	74%	31%	48%	Rejected
Request by surgeon	78%	22%	38%							Rejected
Patient position during regional anesthesia procedure	74%	69%	71%	91%	82%	84%				Accepted
Monitors applied	91%	78%	82%	95%	75%	81%				Accepted
Baseline vital signs	91%	69%	76%	100%	71%	79%				Accepted
Baseline visual analogue score	35%	33%	33%							Rejected
Pre-medication (type and quantity of sedation)	91%	89%	90%	95%	93%	94%				Accepted
Level of sedation (no sedation / light sedation / deep sedation / general anesthesia)	78%	80%	79%	73%	80%	78%				Accepted
Rationale for block performance under spinal, epidural or general anesthesia	48%	35%	38%							Rejected
Time and date of regional anesthesia procedure	100%	96%	97%	100%	100%	100%				Accepted
Aseptic agent used	100%	70%	79%	100%	76%	83%				Accepted
Sterile gloves used (if applicable)	83%	59%	66%	95%	64%	73%	5%	7%	6%	Rejected
Mask used (if applicable)	83%	54%	62%	77%	60%	65%	5%	7%	6%	Rejected

Gown used (if applicable)	35%	50%	45%							Rejected
Ultrasound probe decontaminated according to local requirements	39%	43%	42%							Rejected
Sterile ultrasound transducer cover used (if applicable)	74%	56%	61%	82%	65%	70%	5%	0%	2%	Rejected
Sterile drape used (if applicable)	70%	54%	58%	86%	53%	62%	0%	0%	0%	Rejected
Aseptic technique used as per local policy							84%	69%	75%	Accepted
Skin infiltration with local anesthetic	96%	57%	69%	86%	80%	82%				Accepted
Needle design: tip, manufacturer, length, gauge	100%	87%	91%	95%	91%	92%				Accepted
Local anesthetic used for regional anesthesia technique (concentration and volume)	100%	100%	100%	100%	100%	100%				Accepted
Adrenaline dose if used (concentration)	100%	94%	96%	100%	98%	99%				Accepted
Adjunct / additive used (bicarbonate, clonidine etc.)	100%	100%	100%	100%	100%	100%				Accepted
Side of block	100%	100%	100%	100%	100%	100%				Accepted
Blocked side marked	91%	65%	73%	82%	62%	68%	32%	25%	28%	Rejected
Technique of needle localization (ultrasound / nerve stimulator / landmark)	100%	100%	100%	100%	100%	100%				Accepted
Approach used (in plane / out of plane)	57%	80%	73%	64%	75%	71%	22%	17%	19%	Rejected
No Evoked Motor Response < ___mA (when applicable, e.g. when nerve stimulator used)	74%	70%	71%	86%	80%	82%				Accepted

Minimum current and current duration (if nerve stimulator used)	43%	67%	60%	82%	84%	83%				Accepted
Description of motor response (if nerve stimulator used)	65%	48%	53%	91%	65%	73%	22%	14%	17%	Rejected
Description of quality of paresthesia (if using paresthesia technique)	70%	46%	53%	86%	55%	64%	11%	3%	6%	Rejected
Catheter tip location confirmed by ultrasound / nerve stimulator	57%	70%	66%	59%	62%	61%	0%	7%	4%	Rejected
Technique of injection (via needle and/or catheter)	52%	63%	60%	68%	60%	62%	16%	17%	17%	Rejected
Note on incremental injection	83%	48%	58%	91%	53%	64%	21%	0%	8%	Rejected
Extra-neural spread visualized	61%	65%	64%	82%	67%	71%	16%	24%	21%	Rejected
Presence / absence nerve swelling							11%	24%	19%	Rejected
Note on resistance during injection (pressure < 15psi)	52%	57%	56%	59%	60%	60%	6%	10%	8%	Rejected
Absence of blood on aspiration	96%	81%	86%	95%	84%	87%				Accepted
Needle depth before injection	57%	33%	40%							Rejected
Catheter depth at the skin	83%	91%	88%	82%	95%	91%				Accepted
Absence of pain / paresthesia during injection	100%	85%	90%	91%	85%	87%				Accepted
Technique (approach used e.g. median / paramedian)	96%	93%	94%	100%	93%	95%				Accepted
Vertebral level of needle insertion	100%	98%	99%	100%	98%	99%				Accepted
Technique used: Loss of resistance to saline/air for epidural insertion	91%	93%	92%	100%	96%	97%				Accepted
Number of attempts	83%	93%	90%	95%	93%	94%				Accepted

Epidural needle depth at loss of resistance	96%	96%	96%	91%	96%	95%				Accepted
Catheter depth at the skin	96%	98%	97%	95%	96%	96%				Accepted
Method used to secure catheter				59%	40%	45%				Rejected
Note on aspiration and action taken	100%	87%	91%	100%	82%	87%				Accepted
Epidural test dose (if applicable)	100%	93%	95%	100%	96%	97%				Accepted
Absence of pain / paresthesia during injection	96%	85%	88%	86%	84%	84%				Accepted
Dermatomal level of spinal or epidural block achieved (if assessed)	65%	80%	75%	100%	87%	91%				Accepted
Complications (neuraxial)	100%	98%	99%	100%	100%	100%				Accepted
Adequacy of regional anesthesia procedure	74%	69%	70%	59%	75%	70%	22%	45%	36%	Rejected
Complications (nerve block)	100%	96%	97%	100%	98%	99%				Accepted
Ultrasound image included in patient record (if US used)				82%	25%	42%				Rejected
Patient vital signs after the procedure	87%	65%	71%	95%	71%	78%				Accepted
Post block monitoring completed by _____/ handed over/off to _____(e.g. Recovery, PACU)				73%	45%	53%	17%	0%	6%	Rejected
Patient visual analogue score after the procedure	39%	41%	40%							Rejected
Post procedure instructions	65%	91%	83%	73%	89%	84%				Accepted