



OPEN ACCESS

Factors associated with use of opioid rescue medication after surgery

Isabell M Meier ,¹ Marie Eikemo ,^{1,2} Martin Trøstheim ,^{1,2} Kaja Buen,³ Eira Jensen,³ Siri Gurandsrud Karlsen,³ Silje E Reme ,^{2,4} Chantal Berna ,⁵ Siri Leknes ,^{1,2} Gernot Ernst ,^{2,3}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rapm-2023-104412>).

¹Department of Physics and Computational Radiology, Oslo University Hospital, Oslo, Norway

²Department of Psychology, University of Oslo, Oslo, Norway

³Department of Anesthesiology, Kongsberg Hospital, Vestre Viken Hospital Trust, Kongsberg, Norway

⁴Department of Pain Management and Research, Oslo University Hospital, Oslo, Norway

⁵Centre of Integrative and Complementary Medicine, Division of Anaesthesiology, Lausanne University Hospital, Lausanne, Switzerland

Correspondence to

Dr Isabell M Meier, Department of Physics and Computational Radiology, Oslo University Hospital, Oslo 0372, Norway; i.m.meier@psykologi.uio.no

SL and GE contributed equally.

Received 27 February 2023

Accepted 23 June 2023



© American Society of Regional Anesthesia & Pain Medicine 2023. Re-use permitted under CC BY-NC. No commercial re-use. Published by BMJ.

To cite: Meier IM, Eikemo M, Trøstheim M, et al. *Reg Anesth Pain Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/rapm-2023-104412

ABSTRACT

Background Opioid exposure after surgery increases risk of persistent opioid use. Here, we characterize at-home use of opioid rescue medication during 1–2 days after outpatient surgery (N=270) in a postoperative opioid-sparing context at a Norwegian hospital.

Methods The postsurgical pain management plan included non-steroidal anti-inflammatory drugs and up to six pills of 5 mg oxycodone as rescue analgesics. In this observational study we assessed risk factors for taking rescue opioids after surgery, by comparing patients who did, with those who did not.

Results Only 35% (N=228) of patients reported taking rescue opioids 1–2 days after discharge. Patients taking rescue opioids after surgery (opioid-takers) differed from non-takers by prevalence of preoperative chronic pain (>3 months; 74% vs 48%), higher pain severity and interference before and after surgery, reporting lower ability to cope with postsurgical pain, higher nervousness about the surgery, being younger, and having received more opioid analgesics in the recovery room. Exploratory predictive modeling identified opioid administration in the recovery room as the most important predictor of at-home rescue medication use. Follow-up after >4 months indicated low acute pain levels (mean±SD = 1.1±1.8), with only four patients (2%, N=217) reporting opioid analgesic use.

Conclusion Factors related to at-home rescue medication use closely mirrored known risk factors for persistent opioid use after surgery, such as prior chronic pain, prior substance use, affective disturbances, and pain severity before surgery. These findings are potential targets in patient-centered care. Nevertheless, and reassuringly, findings are consistent with the idea that opioid-sparing postsurgical care can prevent large-scale chronic opioid use.

INTRODUCTION

Opioid analgesics are a cornerstone of postsurgical pain treatment. Prolonged opioid use and large prescription sizes are associated with heightened prevalence of opioid misuse, addiction and mortality.^{1–3} Over-prescription was a root cause for the first wave of the opioid epidemic in North America⁴; outside of the USA and Canada opioid-sparing postoperative analgesic regimens are more commonly chosen to prevent persistent opioid use. Overall, studies on acute postoperative opioid use and opioid sparing pathways are conducted in the North American context,^{5–8} where opioid

SWHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Growing evidence indicates that risk factors for persistent opioid use include acute and chronic pain, history of substance use, younger age and affective disorders.

WHAT THIS STUDY ADDS

⇒ The study demonstrates strong overlap between risk factors for persistent opioid use and acute use of opioids analgesics in an opioid sparing setting and highlights acute pain before surgery and frequency of opioid administration in the recovery room as relevant predictors for opioid rescue medication use.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The study demonstrates the feasibility of opioid sparing postoperative pain management after outpatient surgery and suggests potential targets for early, preventive patient-centered care.

prescriptions as well as opioid prescription fillings are largely higher than internationally and opioid sparing prescription practice is scarce.^{9 10}

In opioid-sparing postoperative care, non-opioid analgesics are the main treatment and limited opioid doses are provided to be taken as needed (ie, ‘rescue medications’). In contrast to the growing evidence on predictors of persistent opioid use, which include history of substance use and chronic pain, affective disorders and pain severity,^{1 11–13} little is currently known about which patient characteristics relate to postsurgical opioid use during the acute recovery phase when patients self-administer analgesics from home. Initial data on postoperative opioid use from home in a *voluntary* opioid-sparing postoperative pathway indicated that opioid-takers reported higher pain levels after surgery, lower coping ability and were on average younger than patients who did not take opioid rescue medication.⁷

Here, we used prospective observational data to investigate at-home use of opioid rescue medication after outpatient surgery at a Norwegian hospital, where postoperative opioid-sparing treatment is standard care. Patients who either did (N=81) or did not (N=147) self-administer opioid analgesics in the first 1–2 days after surgery were compared with the aim of identifying potential predictive




Figure 1 Inclusion flow chart. Of the 270 patients included in the study, 228 were reached for a phone interview during acute recovery (1–2 days after surgery). Of the 230 patients reached for long-term follow-up (4–30 months after surgery), 218 consented to take part in the phone interview and to receive questionnaires. From 186 of those 218 patients we also have information on postoperative opioid use during acute recovery. 151 of the 218 patients filled in at least one questionnaire.

measures of postoperative rescue opioid use. Since opioid use during acute recovery can be a first step in the development of persistent opioid use, we hypothesized that risk factors for persistent opioid use (eg, younger age, higher acute pain, history of chronic pain, prolonged opioid use) would also be related to postoperative use of rescue opioids in the 2 days after surgery (acute recovery phase). Unlike previous studies,^{7,8} our prospective study assessed use of postoperative opioid rescue analgesics in a context where opioid sparing postoperative pain management is standard of care.

METHODS

Participants and procedure

A convenience sample of patients (N=270, 153 women, age mean±SD = 47.5±14.2) scheduled for outpatient surgery with general anesthesia was recruited at Kongsberg Hospital in Norway between April 2018 and June 2021. The participants

were adult patients healthy enough to undergo minor outpatient surgery, classified as ASA-I–II (American Society of Anesthesiologists)¹⁴ (see online supplemental file). Participants were invited to join the study and signed written informed consent on arrival at the hospital. Contact attempts for follow-up measures were limited to three times for phone interviews and three email reminders for online questionnaires resulting in varying response rates and a potential response bias. An inclusion flowchart for the different time points reported can be found in figure 1.

Participants received treatment as usual, that is, according to hospital routines. At discharge from the hospital after surgery, patients received two types of analgesics to take home: A fixed combination of naproxen (500 mg) with esomeprazole (20 mg) and as rescue medication up to six oxycodone pills (5 mg). The number of oxycodone pills was decided by the medical personnel, considering the likelihood of high pain and the accessibility of a pharmacy/general practitioner. Patients were instructed to rely on non-opioid analgesics as much as possible for pain management and to take one initial dose of paracetamol (1.5 g for <75 kg body weight and 2g for >75 kg body weight) followed by 1 g of paracetamol up to four times a day, non-steroidal anti-inflammatory drug if no contraindications were present, and opioids only if absolutely necessary. A detailed overview of the perioperative pain management can be found in the online supplemental material.

Clinical and research specific data were collected over six time points: weeks before surgery, ~60 min before surgery, on the surgery table, in the recovery room, during acute recovery (1–2 days after discharge) and at long-term follow-up (4–30 months after surgery) (figure 2). Data on acute responses to opioids on the operating-table are reported elsewhere.¹⁵

Outcomes

The **grouping variable self-reported opioid use during acute recovery** was collected during the phone interview in the acute recovery phase (1–2 days after discharge) and defined as having taken (1) at least one pill of the oxycodone tablets provided by the hospital or (2) any other opioid analgesic patients had available. **Patient characteristics** were collected via self-report on paper questionnaires and the hospital database and include *sex, age, body mass index (BMI), operation category, opioid received on the surgery table, tobacco use, history of pain, history of*




Figure 2 Timeline of data collection. 2–3 weeks prior to surgery: As part of the hospital routine, patients filled in a form on demographics, levels of nervousness before surgery, pain and pain interference. In the hour before surgery, patients completed questionnaires on mood, pain, pain interference and previous experience with opioids. In the operating room, patients rated their mood and acute drug effects verbally. Recovery room: Patients received opioid or non-opioid analgesics up to five times in the recovery room. Data was extracted from the hospital system. Acute recovery: 1–2 days after surgery. In a semi-structured phone interview, patients reported on their use of the analgesics, rated pain intensity, interference and coping as well as mood and satisfaction with the hospital treatment and stay. Follow-up: 4–30 months after surgery. In a second brief phone interview, patients reported on the extended recovery period after surgery (pain control, social support), current medication use and work status. We also included questionnaires on their substance use, experience of life stressors, pain and pain catastrophizing, life satisfaction, hedonic experience and mindfulness in day-to-day life. Long-term follow-up was an add-on to the project, leading to variability in the time since surgery (4–30 months).

opioid use and analgesic use in the recovery room (number of doses administered). Subjective ratings (0–10 rating scales) of pain indices and nervousness were collected via self-report on paper questionnaires in the weeks before surgery and on day of surgery and via phone interview during acute recovery and long-term follow-up; other state measures (feeling good, feeling anxious) were measured verbally on the operating table before and after induction of the opioid analgesic. Patient satisfaction was collected in the phone interview during acute recovery.

Subjective ratings

Patients rated their pain intensity and pain interference, nervousness about the surgery or recovery at three time points: weeks before surgery, morning of surgery and during the acute recovery phase 1–2 days after discharge from the hospital. Ratings of feeling good and anxious were collected on the surgery table before and after opioid injection. Patient satisfaction was measured during the acute recovery phase 1–2 days after discharge from the hospital. Details can be found in the online supplemental material.

Long-term follow-up questionnaire data

We measured pain¹⁶ and pain catastrophizing,^{17 18} depression and anxiety,¹⁹ childhood trauma,^{20 21} subjective perception of socio-economic status during childhood and adulthood,²² potential substance abuse,^{23 24} general life satisfaction,²⁵ hedonic experience,²⁶ and mindfulness in day-to-day life.²⁷ An overview of all questionnaires can be found in the online supplemental material.

Statistical analyses

All analyses were performed in R V.4.1.2.²⁸

Primary analysis

To determine the relevance of factors previously associated with *persistent opioid use* (patient characteristics and subjective ratings) in this *acute recovery context* we compared patients who either did (N=81) or did not (N=147) self-administer opioid analgesics within the first 2 days after surgery using non-parametric repeated-measures analysis of variance (ANOVA) (rank-based)²⁹ as well as Welch's t-tests, Wilcoxon rank-sum test and χ^2 tests. Additionally, we tested whether the opioid type patients received on the surgery table influenced opioid use in the recovery room and at-home. Since we conducted a large number of pairwise comparisons, we chose a more conservative alpha level of 0.01 to lower the false-positive rate.

Exploratory analyses

To identify important variables for predicting opioid use after surgery, we used multimodel inference³⁰ implemented in R³¹ with the *MuMIn* package.³² Multimodel inference accounts for the uncertainty inherent in stepwise model selection by aggregating information from multiple models.³⁰ Stepwise methods for model selection were preferred historically due to limits in computational power. In contrast, the multimodel approach tests all possible models to determine the general importance (or usefulness) of each individual variable for predicting the outcome of interest. This is done by quantifying the tendency of each individual predictor to appear in good-performing models.

We used binary logistic regression with *opioid use after surgery* as outcome and fitted models for all 2048 possible combinations of a prespecified set of 11 predictors (*sex, age, BMI, previous opioid use, tobacco use, chronic pain, pain weeks before surgery, nervousness weeks before surgery, operation category, opioid*

administration in the recovery room, and pain on the day of surgery), excluding interactions.

Akaike weights were calculated from Akaike information criteria (AIC) for all models to indicate the probability of each model being the best model.³³ The importance of each predictor across models was determined by calculating the sum of Akaike weights of all models containing the predictor.³⁰ To assess the overall significance of each predictor, we calculated the full-average model coefficients across all models.^{30 34}

Due to missing data, we performed multimodel inference on the complete cases (n=118) and on all cases after imputation (n=228). Data for a total of 48% of incomplete cases were imputed. The percentage of imputed data varied by predictor with the lowest amount of imputed data for tobacco use (2%) to the highest amount of imputed data for chronic pain (33%). Multiple imputation by chained equations was implemented with the *mice* package.³⁵ We used a quadratic model³⁶ implemented with the *howManyImputations* package³⁷ to determine the required minimum number of 48 imputations for the analysis. All imputations were iterated 10 times.³⁸ We used the *mami* package^{39 40} to pool multimodel inference results across imputations.

RESULTS

Patient characteristics

Descriptive data of the full patient population are available in [table 1](#). Approximately half of the participants reported previous pain that lasted more than 3 months (56% of N=180). Previous experience with opioids was limited: Of 242 patients 34% reported no prior opioid experience (opioid-naïve), 60% had taken an opioid analgesic at least once. Only 6% of N=232 reported to have used opioids for more than 2 weeks at a time. In the interview 1–2 days after surgery (N=228), most patients (99%) reported to be satisfied with their treatment and stay at the hospital.

Analgesic use after surgery

Recovery room

Of the 270 patients, 200 (74%) received an analgesic (opioid or non-opioid) in the recovery room. Of these, 110 patients (55%) received at least one opioid dose. Only 54 (49%) of these 110 patients received more than one opioid analgesic dose. We found no difference of the opioid administration in the recovery room depending on the history of opioid use. Among the patients who provided information on their opioid history (N=242), 39% of opioid-naïve patients received opioids in the recovery room, compared with 40% of patients with a history of opioid use ($\chi^2=0.01(1)$, $p=0.92$). A larger proportion of patients receiving preoperative oxycodone compared with remifentanyl received opioids in the recovery room (note that preoperative opioids were not randomized). Detailed results can be found in the online supplemental material.

Acute recovery at home

Most patients (93.2%, N=265) received one to six pills of 5 mg oxycodone as rescue medication (median=5, IQR=2–5). Only 35.5% of patients reached at 1–2 days postsurgery (N=228) reported use of oxycodone (32%) or another opioid analgesic they already had at home (3.5%).

Long-term analgesic use

In the long-term follow-up phone interview, 24 (11%) of the patients reaching and consenting to take part (N=218) reported

Table 1 Overview of patient characteristics for the full sample and the two subsamples

	Full sample	Opioid-takers	Non-opioid-takers	P value
N	270	81	147	
Sex, n (%)	153 women (57)	52 women (64)	80 women (54)	0.2*
Age, mean (SD)	47.5 (14.2)	44.5 (13.4)	49.7 (13.8)	0.006†
BMI, mean (SD)	27 (5.1)	28.4 (6.2)	26.5 (4.3)	0.3‡
Surgery type, n (%)				
Colorectal	48 (18)	9 (11)	26 (18)	0.26*
Minor abdominal	108 (40)	36 (44)	54 (37)	0.3*
Minor gynecological	59 (22)	14 (1)	41 (28)	0.1*
Otorhinolaryngology	6 (2)	2 (3)	2 (1)	0.9*
Minor orthopedics	32 (12)	14 (17)	15 (10)	0.2*
Other	17 (6)	6 (7)	9 (6)	0.9*
Analgesics received in the recovery room, n (%)				
No analgesic	70 (26)	13 (16)	48 (33)	<0.001*
Non-opioid analgesic	89 (33)	14 (17)	61 (42)	
Opioid analgesic	111 (41)	54 (67)	38 (26)	
Prior pain, >3 months, n (%)	101 (56)	36 (72)	49 (48)	0.007*
Tobacco use (yes/no), n (%)	46 (17)	15 (19)	21 (15)	0.5*
History of opioid exposure				
Opioid-naive, n (%)	82 (34)	20 (27)	52 (41)	0.07*
Opioid use >2 weeks, n (%)	14 (6)	5 (7)	5 (4)	0.5*
Satisfied w/treatment, n (%)	223 (99)	78 (96)	145 (99)	0.6*
Opioid use at long-term follow-up, n (%)	4 (2)	4 (6)	0	0.2*

Total counts (n), percentages (%) and averages (mean) for the full sample (n=270) and per group for the patients reached during acute recovery (n=228). Opioid use at long-term follow-up was measured in the subsample of n=218 who consented to take part in the follow-up part of the study. Percentage of missing data for group comparisons: pain >3 months: 33%; tobacco: 2%; previous opioid use: 11%; opioid use >2 weeks: 15%; opioid use long-term follow-up: 19%. P values reflect comparisons of opioid-takers vs non-takers.
 †Bold p values indicate p<.01.
 * χ^2 test.
 †Welch's t-test.
 ‡Wilcoxon rank-sum test.
 BMI, body mass index.

current pain medication use, of which only 4 patients reported opioid analgesic use (1.8%).

Comparison of patients who did and did not use rescue opioids during acute recovery

Characteristics of opioid-takers and non-takers are listed in [table 1](#). Results on acute opioid effects on the surgery table can be found in online supplemental material.

Pain and pain coping

A significantly higher percentage of the opioid-takers reported having had pain for more than 3 months before surgery (72%) compared with the non-takers (48%, N=228, likelihood-ratio chi-square (χ^2)=7.2, p=0.007). Opioid-takers were also more likely to have received opioids in the recovery room (N=228; 67% vs 26%, (χ^2)=36.3, p<0.001).

The non-parametric repeated-measures ANOVA on pain intensity ratings (N=228) revealed a significant main effect of group (opioid-takers vs non-takers; $F(1)=36.5$, p<0.0001) and time (weeks before surgery, day of surgery, acute recovery, long-term follow-up; $F(2.8)=68.9$, p<0.0001). No significant group×time interaction effect was found ($F(2.8)=2.0$, p=0.12). As illustrated in [figure 3](#), opioid-takers reported significantly higher pain intensity weeks before surgery, on the day of surgery and during acute recovery, but not at long-term follow-up when low levels of current pain were reported (online supplemental table 2).

The pattern of pain interference ratings strongly resemble pain intensity ratings across time and groups (online supplemental table 2).

Wilcoxon rank-sum tests showed that during acute recovery, both groups reported experiencing high pain relief from the analgesic medications ingested (N=210, mean±SD: opioid-takers:

6.8±2.3; non-takers: 7.1±2.8; W=5719, p=0.2). While pain coping was rated high overall, opioid-takers' ratings were significantly lower (N=225, opioid-takers: 7.9±2.0; non-takers: 8.9±1.5; W=7778, p<0.001).

Nervousness

A non-parametric repeated-measures ANOVA revealed a significant main effect of group (N=228, opioid taker vs non-taker; $F(1)=8.3$, p=0.004) and time (weeks before surgery, on the day of surgery and acute recovery; $F(1.8)=69.3$, p<0.0001),

Pain intensity ratings across time




Figure 3 Pain intensity ratings for opioid-takers and non-takers. Opioid-takers reported significantly higher pain intensity before surgery and during acute recovery compared with non-takers. The plots display groupwise means, 95% CIs, dots depict individual numeric ratings indicated via questionnaire (weeks before surgery, day of surgery, long-term follow-up) and verbally via phone interview (acute recovery). Numeric ratings were always given as discrete numbers on a scale from 0 to 10. Error bars represent 95% CIs. ***p≤0.001; ns, not significant.




Figure 4 Nervousness ratings for opioid-takers and non-takers. Opioid-takers were significantly more nervous about the surgery in the weeks before but did not differ significantly from non-takers in nervousness on the day of surgery or about recovery in the 2 days after surgery. Groupwise means, 95% CIs, dots depict individual numeric ratings indicated via questionnaire (nervousness about surgery, weeks before and day of surgery) and verbally via phone interview (nervousness about recovery, 1–2 days after surgery). Numeric ratings were always given as discrete numbers on a scale from 0 to 10. Error bars represent 95% CIs. ** $p < 0.01$, ns, not significant.

however no significant interaction effect ($F(1.8) = 1.6$, $p = 0.20$). Opioid-takers rated on average significantly greater nervousness about the surgery weeks before surgery, but not in the hour before surgery and in the acute recovery interview (figure 4).

Long-term follow-up questionnaire data

For questionnaires collected in the 4–30 months after surgery reporting on early trauma, alcohol misuse, drug misuse, pain catastrophizing, pain intensity, depression and anxiety, life satisfaction, perceived socioeconomic status during childhood and adulthood, anhedonia, and mindfulness, no significant differences were found between opioid-takers and non-takers (online supplemental table 3).

Predictors of acute opioid use after surgery

Opioid administration in the recovery room was identified as the most important and only significant predictor of opioid use after surgery in the averaged models using multimodel inference, when analyzing both complete cases ($N = 118$, ORs for complete cases (OR_{comp}) (95% CI) = 2.89 (1.70 to 4.93), $p_z < 0.0001$, $Importance = 1$) and all cases after imputation ($N = 228$, ORs for imputed cases (OR_{imp}) (95% CI) = 2.68 (1.86 to 3.86), $p_z < 0.0001$, $Importance = 1$; figure 5; online supplemental table 4). A forest plot including model-averaged ORs and indicators of importance for each predictor per model can be found in figure 5. Overall, the pattern of results was similar for multimodel analysis of complete cases and the 48 imputed data sets. Although not significant, the second most important predictor of opioid use after surgery was pain in the weeks before surgery (OR_{imp} (95% CI) = 1.23 (0.98 to 1.54), $p_z = 0.07$, $Importance = 0.88$), followed by age, BMI and history of opioid use ($Importance$ imputed cases: > 0.5 ; see online supplemental table 4).

DISCUSSION

Here, we described opioid rescue medication use in 270 outpatient surgery patients in an opioid-sparing postoperative context. Opioid-sparing postoperative analgesic regimens aim to prevent persistent opioid use. Patients received opioids immediately

Multimodel inference




Figure 5 Identification of important predictors of opioid use after surgery through multimodel inference (2048 possible models). Results based on complete cases before imputation ($n = 118$) are displayed in black while results based on all cases after imputation ($n = 228$) are displayed in gray. As a reference category, we used *Men* for the predictor *Sex*, *No* for the predictors *Tobacco use* and *Chronic pain*, and *Minor abdominal surgery* for the predictor *Operation category* (*Op. cat.*). Black dots and horizontal lines indicate model-averaged ORs and 95% CIs, respectively. Arrows indicate that the 95% CI exceeds the limits of the x-axis. ORs > 1 indicate increased risk of opioid use after surgery while ORs < 1 indicate reduced risk of opioid use after surgery. Horizontal bars indicate the importance of each predictor, calculated as the sum of Akaike weights for models containing the predictor. Models containing predictors of high importance tend to perform better at predicting opioid use after surgery compared with models not containing these predictors.

before and during surgery (remifentanyl during anesthesia), but only 41% received opioids in the recovery room. While most patients (93.2%) received one to six pills of 5 mg oxycodone as rescue medication for at-home postoperative pain management, 65% of the interviewed patients had not taken any opioid analgesic 1–2 days after discharge. Since the risk of persistent use hinges on initial use, we compared the rescue opioid-takers to the non-takers, focusing on known risk factors for persistent postsurgical opioid use. Consistent with these known risk factors, we found that opioid-takers were of significantly younger age, more likely to have chronic pain, had higher levels of pain before surgery and during acute recovery, lower pain coping ability after surgery, higher nervousness about the surgery in the weeks before, and were more likely to have received opioids in the recovery room.

Exploratory predictive modeling identified frequency of opioids received in the recovery room after surgery as the most important predictor for at-home use of opioid rescue medication, pointing to postoperative pain management in the recovery room as a relevant target for preventive care. Other factors identified as important in a large proportion of models tested, though not significant in the average model, included pain severity in the weeks before surgery, age, BMI, and history of opioid use. Nevertheless, only 2% of patients were still on opioids at long-term follow-up, supporting the idea of efficient prevention through the postoperative opioid-sparing scheme.

Compared with a Northern-American non-opioid-sparing context, where over 75% of patients fill opioid prescriptions after surgery^{2,10} and 4.5% take opioids over 3 months or longer,¹ the current patient population had very little opioid exposure.

Patient endorsement of treatment satisfaction was high in both groups. This is consistent with findings from studies introducing the opioid-sparing approach in the USA.^{7,8} The overall pain level at long-term follow-up was low and significantly reduced in comparison to presurgery and acutely postsurgery. Taken together, the high satisfaction, low levels of persistent pain and analgesic use months after surgery, validate the opioid-sparing approach as an efficient pain management scheme within this population, in line with the idea that opioid-sparing post-surgical care can prevent large-scale chronic opioid use.

The aim of this study was to identify whether known risk factors of persistent opioid use can also explain rescue opioid use after surgery. In line with previous studies indicating that elevated levels of acute and chronic pain are major vulnerability factors for persistent opioid use,^{11,41} we found that patients who self-administered opioids during acute recovery (opioid-takers) were more likely to have chronic pain, and reported higher levels of acute pain before surgery, on the day of surgery, and in the 2 days after surgery. Consistent with their elevated pain ratings, opioid-takers were also more likely to receive opioids in the recovery room. While the recovery room staff were not blinded to the patients' opioid history which can result in a potential bias, we found no significant association between opioid history and opioid administration in the recovery room.

Both frequency of opioid administration in the recovery room and pain severity in the weeks before surgery emerged as relevant predictors for at-home use of opioid analgesics, making them interesting targets for preventive care. Patients who took opioids after surgery also reported to cope less well with postsurgical pain, although the overall reported pain relief in the two groups was comparable. This is in line with research demonstrating that improving pain coping skills significantly reduced opioid craving in adults with chronic pain and opioid dependency,⁴² and postoperative opioid use in surgery patients.⁶

Age has previously been identified as an important predictor of postoperative opioid requirements, with younger patients requesting substantially higher doses of morphine.⁴³ In line with these findings, although the age range for the two groups was similar, opioid-takers were on average 5 years younger than non-takers. Opioid-takers and non-takers did not differ significantly in sex, BMI and history of tobacco use. All patients were selected for minor ambulatory surgery, hence excluding those with massive noxious input such as thoracotomy or major abdominal surgery. Despite potential systematic differences in the noxious input of the surgery, there was no significant difference in acute opioid use depending on the surgery type, nor was surgery type a significant predictor of postoperative opioid use in our sample. This is in line with prior evidence that new persistent opioid use and postoperative opioid use in the 2 months after surgery was not dependent on the type or severity of surgery patients received (eg, minor vs major surgery).^{6,7,11}

Undergoing surgery is considered a major stressor and anxiety is reported as the worst aspect of the perioperative experience.⁴⁴ Opioid-takers were on average significantly more nervous about the surgery weeks before compared with non-takers. When asked how nervous they felt on the day of surgery and later about their recovery in the postoperative phone interview, the opioid-takers and non-takers reported comparably low numbers. We speculate that the limited nervousness in the postoperative period could be a consequence of their positive experience at the hospital, as documented by the overall high satisfaction rate.

Anxiety, depression^{11,41,45} and catastrophizing^{12,13,46} have been reported to be independently associated with persistent opioid use after surgery, unrelated to factors such as severity of injury,

treating surgeon or surgery type.¹³ We found no differences in anxiety, depression or catastrophizing scores in our long-term follow-up data, however it is possible that differences were present before surgery and/or during acute recovery. We found no significant differences between opioid-takers and non-takers in history of trauma, perceived socioeconomic status, or experience of loneliness.

LIMITATIONS

The study had several limitations. While the purely observational data ensure high ecological validity, relying on self-reports of rescue opioid use in the postsurgical phone interview could result in bias by desirability effects. Some questionnaires were only administered months or years after surgery, limiting the ability to interpret any lack of association with opioid self-administration. The low overall opioid exposure of the sample limits our ability to test the relationship between prior opioid use and opioid taking in the days after surgery. Similarly, some of the comparisons between subgroups were underpowered (eg, opioid use at long-term follow-up). Since we aimed for a descriptive approach, we conducted a high number of statistical tests with no correction for multiple comparison beyond a more stringent alpha level ($\alpha=0.01$). We tested for the predictive value of several factors collected before surgery using a multimodel approach with logistic regressions. While our findings align with and add to previous retrospective data,⁸ we recommend that our findings should be replicated.

CONCLUSION

With opioid exposure being a large risk factor for persistent use, we need to understand what predicts the intake of rescue medication. Overall, only 36% of patients reported opioid analgesic use within 1–2 days after surgery. Opioid-takers were younger, more likely to have chronic pain and reported higher nervousness before surgery as well as higher pain levels and worse pain coping skills overall. Pain severity before surgery and frequency of opioids administered before discharge are relevant predictors of opioid use after surgery in an opioid-sparing context and could be targets for patient-centered care. The data reported here demonstrate the feasibility and utility of the opioid-sparing postsurgical care, with high patient satisfaction throughout, adequate postoperative pain control and very low incidence of persistent opioid analgesic use.

Twitter Isabell M Meier @IM_Meier, Marie Eikemo @Marie_Eikemo, Martin Trøstheim @martintrostheim, Chantal Berna @Chantal_berna_, Siri Leknes @sirileknes and Gernot Ernst @gernot_ernst

Contributors Study conception and design: SL, ME, GE, IMM. Acquisition of data: GE, KB, EJ. Analysis and interpretation of the data: IMM, MT, SGK, ME, SL. Drafting of manuscript: IMM, ME, MT, SL, CB, SGK. Critical revision: CB, SER, GE. SL is the guarantor for the data, ensuring that questions related to the accuracy of the manuscript are appropriately investigated and resolved.

Funding This work was supported by the European Research Council under the EU Horizon 2020 research and innovation programme (grant agreement no. 802885) to SL. IMM was supported by South-Eastern Norway Regional Health Authority (grant: 2020087) to SL, ME and GE.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Procedures were approved by the local data protection officer and Regional Ethics Committee (Rek Sør-Øst D:198224). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. The data has not been made publicly available due to the sensitive nature of health information. The conditions of

our ethical approval, and the constraints of Norwegian law, prevent sharing of the data supporting this research with any individual outside the author team.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, an indication of whether changes were made, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Isabell M Meier <http://orcid.org/0000-0002-7606-1059>
 Marie Eikemo <http://orcid.org/0000-0001-5103-919X>
 Martin Trøstheim <http://orcid.org/0000-0002-6021-6714>
 Silje E Reme <http://orcid.org/0000-0001-5870-4906>
 Chantal Berna <http://orcid.org/0000-0001-8258-7412>
 Siri Leknes <http://orcid.org/0000-0002-6667-4601>
 Gernot Ernst <http://orcid.org/0000-0001-5224-0338>

REFERENCES

- Agarwal S, Shah A, Gunaseelan V, et al. New persistent opioid use after surgery in patients with a history of remote opioid use. *Surgery* 2022;171:1635–41.
- Howard R, Brown CS, Lai Y-L, et al. Postoperative opioid prescribing and new persistent opioid use: the risk of excessive prescribing. *Ann Surg* 2022. 10.1097/SLA.0000000000005392 [Epub ahead of print 21 Jan 2022].
- Mohamadi A, Chan JJ, Lian J, et al. Risk factors and pooled rate of prolonged opioid use following trauma or surgery: a systematic review and Meta-(Regression) analysis. *J Bone Joint Surg Am* 2018;100:1332–40.
- Ciccarone D. The triple wave epidemic: supply and demand drivers of the US opioid overdose crisis. *Int J Drug Policy* 2019;71:183–8.
- Buono K, Whitcomb E, Guaderrama N, et al. A randomized controlled trial assessing the impact of opioid-specific patient counseling on opioid consumption and disposal after reconstructive pelvic surgery. *Female Pelvic Med Reconstr Surg* 2021;27:151–8.
- Khorfan R, Shallcross ML, Yu B, et al. Preoperative patient education and patient preparedness are associated with less postoperative use of opioids. *Surgery* 2020;167:852–8.
- Hallway A, Vu J, Lee J, et al. Patient satisfaction and pain control using an opioid-sparing postoperative pathway. *J Am Coll Surg* 2019;229:316–22.
- Anderson M, Hallway A, Brummett C, et al. Patient-reported outcomes after opioid-sparing surgery compared with standard of care. *JAMA Surg* 2021;156:286–7.
- Kaafarani HMA, Han K, El Moheb M, et al. Opioids after surgery in the United States versus the rest of the world: the International patterns of opioid prescribing (iPOP) multicenter study. *Ann Surg* 2020;272:879–86.
- Ladha KS, Neuman MD, Broms G, et al. Opioid prescribing after surgery in the United States, Canada, and Sweden. *JAMA Netw Open* 2019;2:e1910734.
- Brummett CM, Waljee JF, Goesling J, et al. New persistent opioid use after minor and major surgical procedures in US adults. *JAMA Surg* 2017;152:e170504.
- Goesling J, Moser SE, Zaidi B, et al. Trends and predictors of opioid use following total knee and total hip arthroplasty. *Pain* 2016;157:1259–65.
- Helmerhorst GTT, Vranceanu A-M, Vrahas M, et al. Risk factors for continued opioid use one to two months after surgery for musculoskeletal trauma. *Journal of Bone and Joint Surgery* 2014;96:495–9.
- Asaa physical status classification system at. n.d. Available: <https://www.asahq.org/standards-and-guidelines/asa-physical-status-classification-system>
- Eikemo M, Meier IM, Løseth GE, et al. Opioid analgesic effects on subjective well-being in the operating theatre. *Anaesthesia* 2023. 10.1111/anae.16069 [Epub ahead of print 28 June 2023].
- Tan G, Jensen MP, Thornby JJ, et al. Validation of the brief pain inventory for chronic Nonmalignant pain. *J Pain* 2004;5:133–7.
- Sullivan MJL, Bishop SR, Pivik J. The pain catastrophizing scale: development and validation. *Psychological Assessment* 1995;7:524–32.
- Fernandes L, Storheim K, Lochting I, et al. Cross-cultural adaptation and validation of the Norwegian pain catastrophizing scale in patients with low back pain. *BMC Musculoskelet Disord* 2012;13:111.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- Bernstein DP, Stein JA, Newcomb MD, et al. Development and validation of a brief screening version of the childhood trauma questionnaire. *Child Abuse & Neglect* 2003;27:169–90.
- Dovran A, Winje D, Overland SN, et al. Psychometric properties of the Norwegian version of the childhood trauma questionnaire in high-risk groups. *Scand J Psychol* 2013;54:286–91.
- Adler NE, Epel ES, Castellazzo G, et al. Relationship of subjective and objective social status with psychological and physiological functioning: preliminary data in healthy white women. *Health Psychol* 2000;19:586–92.
- Babor TF, Higgins-Biddle JC, Saunders JB, et al. *The alcohol use disorders identification test. Guidelines for use in primary care*. World Health Organization, 2001: 41.
- Berman AH, Bergman H, Palmstierna T, et al. *Drug use disorders identification test*. APA PsycTests, 2002.
- Fugl-Meyer AR, Melin R, Fugl-Meyer KS. Life satisfaction in 18- to 64-year-old Swedes: in relation to gender, age, partner and immigrant status. *J Rehabil Med* 2002;34:239–46.
- Snaith RP, Hamilton M, Morley S, et al. A scale for the assessment of Hedonic tone the Snaith-Hamilton pleasure scale. *Br J Psychiatry* 1995;167:99–103.
- Baer RA, Carmody J, Hunsinger M. Weekly change in mindfulness and perceived stress in a mindfulness-based stress reduction program. *J Clin Psychol* 2012;68:755–65.
- R Core Team. R: A language and environment for statistical computing. 2021.
- Noguchi K, Gel YR, Brunner E. nparLD: an R software package for the Nonparametric analysis of longitudinal data in factorial experiments. *J Stat Softw* 2012;50:1–23.
- Burnham KP, Anderson DR. *Formal inference from more than one model: multimodel inference (MMI), model selection and multimodel inference: a practical information-theoretic approach*. New York, NY: Springer New York, 2002: 149–205.
- R Core Team. R: A language and environment for statistical computing. 2022. Available: <https://www.R-project.org>
- Bartoń K. *MuMIn: multi-model inference*. 2022.
- Wagenmakers E-J, Farrell S. AIC model selection using Akaike weights. *Psychon Bull Rev* 2004;11:192–6.
- Symonds MRE, Moussalli A. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behav Ecol Sociobiol* 2011;65:13–21.
- Buuren S van, Groothuis-Oudshoorn K. Mice: multivariate imputation by chained equations in R. *J Stat Soft* 2011;45.
- von Hippel PT. Hippel PT von: how many Imputations do you need? A two-stage calculation using a quadratic rule. *Sociological Methods & Research* 2020;49:699–718.
- Errickson J. howManyImputations: calculate how many Imputations are needed for multiple imputation. 2023. Available: <https://cran.r-project.org/package=howManyImputations>
- White IR, Royston P, Wood AM. Multiple imputation using chained equations: issues and guidance for practice. *Stat Med* 2011;30:377–99.
- Schomaker M. Model averaging and model selection after multiple imputation using the R-package MAMI. 2017. Available: <http://mami.r-forge.r-project.org>
- Schomaker M, Heumann C. Model selection and model averaging after multiple imputation. *Computational Statistics & Data Analysis* 2014;71:758–70.
- Hilliard PE, Waljee J, Moser S, et al. Prevalence of preoperative opioid use and characteristics associated with opioid use among patients presenting for surgery. *JAMA Surg* 2018;153:929–37.
- Messina BG, Worley MJ. Effects of craving on opioid use are attenuated after pain coping counseling in adults with chronic pain and prescription opioid addiction. *J Consult Clin Psychol* 2019;87:918–26.
- Macintyre PE, Jarvis DA. Age is the best predictor of postoperative morphine requirements. *Pain* 1996;64:357–64.
- Walker EMK, Bell M, Cook TM, et al. Patient reported outcome of adult perioperative anaesthesia in the United Kingdom: a cross-sectional observational study. *Br J Anaesth* 2016;117:758–66.
- Nguyen SN, Hassett AL, Hu H-M, et al. Prospective cohort study on the trajectory and association of perioperative anxiety and postoperative opioid-related outcomes. *Reg Anesth Pain Med* 2022;47:637–42.
- Martel MO, Wasan AD, Jamison RN, et al. Catastrophic thinking and increased risk for prescription opioid misuse in patients with chronic pain. *Drug Alcohol Depend* 2013;132:335–41.