	Day 7			Day 14			Day 21			Day 28		
	Class 1	Class 2	Class 3	Class 1	Class 2	Class 3	Class 1	Class 2	Class 3	Class 1	Class 2	Class 3
Number of subjects	322	9	67	291	7	62	249	66	56	215	6	52
				Opioid 1	elated side	effects n(%	)					
Dizziness	73 (23)	0 (0)	24 (36)	48 (16)	1 (14)	15 (24)	31 (12)	1 (17)	13 (23)	11 (5)	0 (0)	6 (11)
Nausea	62 (19)	0 (0)	22 (33)	28 (10)	1 (14)	15 (24) <sup>†</sup>	23 (9)	2 (33)	14 (25) <sup>†</sup>	9 (4)	0 (0)	8 (15)
Vomiting	18 (6)	0 (0)	2 (3)	3 (1)	0 (0)	0 (0)	3 (1)	0 (0)	0 (0)	2 (1)	0 (0)	0 (0)
Constipation	153 (47)	7 (78)	39 (58)	75 (26)	2 (28)	28 (45)	45 (18)	1 (17)	21 (38) <sup>†</sup>	31 (14)	1 (17)	12 (23)
Difficulty urinating	13 (4)	0 (0)	8 (12)	6 (2)	1 (14)	5 (8)	5 (2)	0 (0)	1 (2)	2 (1)	0 (0)	2 (4)
Abdominal discomfort	48 (15)	2 (22)	19 (28)	30 (10)	4 (57)	7 (11)	28 (11)	1 (17)	9 (16)	12 (6)	1 (17)	8 (15)
Drowsiness	154 (48)	5 (56)	43 (64)	96 (33)	3 (43)	30 (48)	59 (24)	1 (17)	22 (39)	37 (17)	3 (50)	18 (35)
Sleeplessness	133 (41)	3 (33)	34 (51)	83 (28)	2 (28)	20 (32)	44 (18)	1 (17)	15 (27)	27 (13)	2 (33)	11 (21)
Pruritus	55 (17)	2 (22)	22 (33)	43 (15)	1 (14)	10 (16)	23 (9)	1 (17)	9 (16)	20 (9)	1 (17)	12 (23)
Mood Changes	34 (11)	1 (11)	11 (16)	31 (11)	0 (0)	7 (11)	18 (7)	1 (17)	6 (11)	12 (6)	0 (0)	8 (15)
			Unsch	eduled cont	act with he	ealth care sy	stem n(%)					1

Supplemental Table 3: Self-reported side effects in the last 7 days at days 7 through 28 of follow-up following joint, bone or spine surgery.

Provider	24 (7)	2 (22)	14 (21) <sup>†</sup>	27 (9)	0 (0)	8 (13)	21 (8)	1 (17)	6 (11)	13 (5)	0 (5)	5 (10)
Emergency room	5 (1)	1 (11)	0 (0)	2 (1)	0 (0)	2 (3)	1 (1)	0 (0)	0 (0)	1 (1)	0 (0)	0 (0)

Data presented as count(% of column).  $\dagger$  = different from Trajectory 1, *P*=0.05 corrected for multiple comparisons.